arXiv:1111.7255v1 [nlin.SI] 30 Nov 2011

The Klein-Gordon Equation and Differential Substitutions of the Form $v = \varphi(u, u_x, u_y)$

M. N. Kuznetsova,[1, a] A. V. Zhiber,[2, b] and A. Pekcan[3, c]

[1] *Ufa State Aviation Technical University*

[2] *Ufa Institute of Mathematics, Russian Academy of Science*

[3] *Istanbul University, Department of Mathematics*

In this paper we present a complete classification of nonlinear differential equations of the form $u_{xy} = f(u, u_x, u_y)$ reduced to the Klein-Gordon equation $v_{xy} = F(v)$ by differential substitutions $v = \varphi(u, u_x, u_y)$, such that $\varphi_{u_x}\varphi_{u_y} \neq 0$.

PACS numbers: 02.30.Jr

Keywords: nonlinear hyperbolic equations, differential substitutions, Klein-Gordon equation

---

[a] Electronic mail: kuznetsova@matem.anrb.ru

[b] Electronic mail: zhiber@mail.ru

[c] Electronic mail: pekcan@istanbul.edu.tr

I. INTRODUCTION

The emergence of the theory of integrable nonlinear partial differential equations is one of the most important developments in mathematical physics over the last quarter of the last century. In this paper, we study nonlinear differential equations of the form

$$u_{xy} = f(u, u_x, u_y). \tag{1}$$

Such equations have applications in many fields of mathematics and physics. First examples of integrable nonlinear hyperbolic equations were discovered in the 19th century in the works of Liouville[1], Bäcklund[2,3], Darboux[4] and other classics[5,6] in the study of surfaces of constant negative curvature. Then the work was long forgotten, and the interest in them revived only in 1970 due to the numerous applications of hyperbolic equations to physical problems (continuum mechanics, quantum field theory, the theory of ferromagnetic materials, etc.), as well as in connection with the development of the inverse scattering method. The number of publications related to hyperbolic integrable equations is so large that we cannot list them all, so we restrict ourselves only to the reviews and some articles, which are close to the subject of our study.

Despite the large number of publications, the classification problem of integrable equations has not yet been completed. For solving a part of this problem we deal with differential substitutions, which play an important role in the study of integrability of differential equations[7–11].

Definition 1 *The relation*

$$v = \Phi\left(u, \frac{\partial u}{\partial x}, ..., \frac{\partial^n u}{\partial x^n}, \frac{\partial u}{\partial y}, ..., \frac{\partial^m u}{\partial y^m}\right)$$

is called differential substitution from equation (1) *in equation*

$$v_{xy} = g(v, v_x, v_y),$$

if the function v *satisfies the above equation for every solution* $u(x, y)$ *of* (1).

One of the effective methods of investigating integrability of nonlinear equations is to use the cascade method of Laplace integration[12] of the linearized equation

$$p_{xy} - f_{u_x} p_x - f_{u_y} p_y - f_u p = 0.$$

This method was proposed by Zhiber, Sokolov, Startsev[13,14] and Anderson, Kamran, Juras[15,16].

The sequence of functions H_i that are uniquely defined by the formulas

$$H_{-1} = -D_y\left(\frac{\partial F}{\partial u_y}\right) + \frac{\partial F}{\partial u_x}\frac{\partial F}{\partial u_y} + \frac{\partial F}{\partial u}, \quad H_0 = -D_x\left(\frac{\partial F}{\partial u_x}\right) + \frac{\partial F}{\partial u_x}\frac{\partial F}{\partial u_y} + \frac{\partial F}{\partial u}$$

$$H_{i+1} = 2H_i - H_{i-1} - (\ln H_i)_{xy}\,,$$

is called a sequence of Laplace invariants and it is base of this method. There is a close relationship between the properties of nonlinear equations (1) and the chain of Laplace invariants. Zhiber and Sokolov[17] gave classification of equations of Liouville type and provided a procedure for finding a general solution by using Laplace invariants. Startsev[18,19] described the properties of the sequence of Laplace invariants of hyperbolic scalar equations, which have substitutions of any number of high-order.

Differential substitutions are special cases of Backlund transformations and allow to obtain solutions of the equations from solutions of known integrable models. It has been shown[20] that nonlinear equations, whose linearizations are related by Laplace transformations of the first and second order, are related by Bäcklund transformations.

There are number of papers dealing with some differential substitutions that are related to nonlinear hyperbolic equations[17–22]. As an example we illustrate the equation[17,23]

$$u_{xy} = S(u)\sqrt{1-u_x^2}\sqrt{1-u_y^2}, \quad \text{where} \quad S'' - 2S^3 + \lambda S = 0$$

and λ is an arbitrary constant. This hyperbolic type equation is reduced to sin-Gordon equation

$$v_{xy} = c \sin v$$

by differential substitution

$$v = \arcsin u_x + \arcsin u_y + P(u), \quad \text{where} \quad P'^2 = 2S' - 2S^2 + \lambda.$$

Therefore, equation

$$u_{xy} = \sin u\sqrt{1-u_x^2}$$

is reduced to sin-Gordon equation by substitution $v = \arcsin u_x + u$ and also equation $u_{xy} = u\sqrt{1-u_x^2}$ by substitution $v = \arcsin u_x$.

Equation

$$u_{xy} = \sqrt{1 - u_x^2}$$

is reduced by $v = \arcsin u_x$ to wave equation $v_{xy} = 0$.

The modified Tzitzeica equation is given by

$$u_{xy} = S(u)b(u_x)\bar{b}(u_y), \quad \text{where} \quad S'' - 2SS' - 4S^3 = 0$$

and the functions b, $\bar{b}$ is determined by the formulae

$$(u_x - b)(b + 2u_x)^2 = 1, \quad (u_y - \bar{b})(\bar{b} + 2u_y)^2 = 1.$$

It is reduced to Tzitzeica equation

$$u_{xy} = c_1 \exp u + c_2 \exp(-2u)$$

by substitution

$$v = -\frac{1}{2}\ln(u_x - b) - \frac{1}{2}\ln(u_y - \bar{b}) + P(u), \quad \text{where} \quad P'^2 - 2SP' - 3S' - 2S^2 = 0.$$

The known Goursat equation

$$u_{xy} = \sqrt{u_x u_y}$$

is reduced by $v = \sqrt{u_x}$ to $v_{xy} = v/4$.

It is known that the hallmark of integrability of the equation is the existence of higher symmetries. In 1979 Zhiber and Shabat[24] have proved that nonlinear Klein-Gordon equation

$$v_{xy} = F(v) \tag{2}$$

had the algebra of higher symmetries if and only if it was equivalent to the Liouville equation

$$v_{xy} = \exp v$$

or to the Tzitzeica equation or to the sin-Gordon equation.

Recently, all the equations of the form (1) reduced to the Klein-Gordon equation (2) by differential substitutions

$$v = \varphi(u, u_x)$$

were described[25].

In this paper we present a complete classification of nonlinear differential equations of the form (1) reduced to the Klein-Gordon equation (2) by differential substitution

$$v = \varphi(u, u_x, u_y), \quad \varphi_{u_x}\varphi_{u_y} \neq 0 \tag{3}$$

and we proved the following theorem.

Theorem 1 *Suppose that equation* (1) *by differential substitution* (3) *is reduced to the Klein-Gordon equation* (2)*. Then the equation* (2) *takes one of the following forms:*

$$v_{xy} = c, \quad v_{xy} = v, \quad v_{xy} = \exp v, \quad v_{xy} = \sin v, \quad v_{xy} = \exp v + \exp(-2v)$$

up to point transformations $v \to \theta(v)$, $x \to \xi(x)$, $y \to \zeta(y)$*. Here* c *is an arbitrary constant.*

II. DIFFERENTIAL SUBSTITUTIONS OF THE FORM $v = \varphi(u, u_x, u_y)$, $\varphi_{u_x}\varphi_{u_y} \neq 0$

Firstly, we note that variables

$$u, \quad u_x, \quad u_y, \quad u_{xx}, \quad u_{yy}, \ldots$$

are independent because we cannot connect them with each other by equation (1) and its differentiated forms. So every mixed derivatives will be expressed by equation (1) and removed from all the expressions. Thus, u_{xy} is replaced by $f(u, u_x, u_y)$, u_{xyy} is replaced by

$$f_u u_y + f_{u_x} f + f_{u_y} u_{yy},$$

and so on.

By substituting (3) in (2) and using the equation (1), we get

$$\begin{aligned} \varphi_u f + u_x\big(\varphi_{uu}u_y + \varphi_{uu_x}f + \varphi_{uu_y}u_{yy}\big) + u_{xx}\big(\varphi_{u_xu}u_y + \varphi_{u_xu_x}f + \varphi_{u_xu_y}u_{yy}\big) + \\ + \varphi_{u_x}\big(f_u u_x + f_{u_x}u_{xx} + f_{u_y}f\big) + \varphi_{u_y}\big(f_u u_y + f_{u_x}f + f_{u_y}u_{yy}\big) + \\ + f\left(\varphi_{u_yu}u_y + \varphi_{u_yu_x}f + \varphi_{u_yu_y}u_{yy}\right) = F(\varphi). \end{aligned} \tag{4}$$

Since the function F depends only on u, u_x and u_y, we should have the terms before u_{xx}, u_{yy} and $u_{xx}u_{yy}$ are equal to zero i.e.

$$u_{xx}u_{yy}: \quad \varphi_{u_xu_y} = 0, \tag{5}$$

$$u_{xx}: \quad \varphi_{uu_x}u_y + \varphi_{u_xu_x}f + \varphi_{u_x}f_{u_x} = 0, \tag{6}$$

$$u_{yy}: \quad \varphi_{uu_y}u_x + \varphi_{u_y}f_{u_y} + f\varphi_{u_yu_y} = 0. \tag{7}$$

Integrating (5) – (7) leads to

$$\varphi = p(u,u_x) + q(u,u_y), \tag{8}$$

$$\varphi_u u_y + \varphi_{u_x} f = A(u,u_y), \tag{9}$$

$$\varphi_u u_x + \varphi_{u_y} f = B(u,u_x). \tag{10}$$

And the remaining terms in (4) give the following equation,

$$f\big(\varphi_u + u_x\varphi_{uu_x} + \varphi_{u_x} f_{u_y} + \varphi_{u_y} f_{u_x} + u_y\varphi_{uu_y}\big) + \varphi_{uu}u_x u_y + \big(u_x\varphi_{u_x} + u_y\varphi_{u_y}\big)\, f_u = F(\varphi). \tag{11}$$

Hence the problem (1) – (3) is reduced to equations (8)–(11). From (9), (10) we get

$$\big(A - u_y\varphi_u\big)\varphi_{u_y} = \big(B - u_x\varphi_u\big)\varphi_{u_x}. \tag{12}$$

We differentiate (12) with respect to u_x, u_y and obtain

$$\big(u_y\varphi_{u_y}\big)_{u_y}\varphi_{uu_x} = \big(u_x\varphi_{u_x}\big)_{u_x}\varphi_{uu_y}. \tag{13}$$

The relation (13) is valid, if one of the following cases holds:

$$\varphi_{uu_x} = 0, \quad \varphi_{uu_y} = 0, \tag{14}$$

$$\varphi_{uu_x} = 0, \quad \big(u_x\varphi_{u_x}\big)_{u_x} = 0, \tag{15}$$

$$\big(u_y\varphi_{u_y}\big)_{u_y} = 0, \quad \varphi_{uu_y} = 0, \tag{16}$$

$$\big(u_y\varphi_{u_y}\big)_{u_y} = 0, \quad \big(u_x\varphi_{u_x}\big)_{u_x} = 0, \tag{17}$$

$$\frac{\big(u_y\varphi_{u_y}\big)_{u_y}}{\varphi_{uu_y}} = \frac{\big(u_x\varphi_{u_x}\big)_{u_x}}{\varphi_{uu_x}} = \lambda(u), \quad \lambda(u) \neq 0. \tag{18}$$

Let us study (18) at first. Using (8), we get

$$\big(u_y q_{u_y}\big)_{u_y} = \lambda(u) q_{uu_y}, \quad \big(u_x p_{u_x}\big)_{u_x} = \lambda(u) p_{uu_x}.$$

Therefore

$$u_y q_{u_y} = \lambda(u) q_u + C(u), \quad u_x p_{u_x} = \lambda(u) p_u + E(u).$$

We integrate these equations and obtain

$$q = \Phi_1\big(u_y\kappa(u)\big) + \epsilon(u), \quad p = \Phi_2\big(u_x\kappa(u)\big) + \mu(u),$$

$$\kappa(u) = \lambda(u)\kappa'(u), \quad \lambda(u)\epsilon'(u) + C(u) = 0, \quad \lambda(u)\mu'(u) + E(u) = 0.$$

It follows that

$$\varphi = \alpha(u) + \Phi_2\big(u_x\kappa(u)\big) + \Phi_1\big(u_y\kappa(u)\big), \tag{19}$$

where $\alpha = \epsilon + \mu$. Further, the point replacement

$$\int \kappa(u)du \to u$$

in (19) gives

$$\varphi = \alpha(u) + \beta(u_x) + \gamma(u_y). \tag{20}$$

Thus, condition (18) provides the form (20). If we combine (14) with (8), we get (20) again.

Let us study (15). Under this case, using (8) yields to

$$p_{uu_x} = 0, \quad \big(u_x p_{u_x}\big)_{u_x} = 0.$$

Hence

$$p = \alpha(u) + c\ln u_x + c_1.$$

Here $c \neq 0$, c_1 are arbitrary constans. Thus, we obtain

$$\varphi = \alpha(u) + c\ln u_x + q(u, u_y).$$

Replacing $\alpha(u) + q(u, u_y)$ by $q(u, u_y)$, we get

$$\varphi = c\ln u_x + q(u, u_y). \tag{21}$$

The condition (16) is symmetrical to (15) under the change of variables $x \leftrightarrow y$.

Under condition (17) with (8) we have

$$\big(u_y q_{u_y}\big)_{u_y} = 0, \quad \big(u_x p_{u_x}\big)_{u_x} = 0.$$

This yields to

$$q = \mu(u)\ln u_y + \epsilon(u), \quad p = \kappa(u)\ln u_x + \delta(u).$$

Consequently, function φ is described by the formula

$$\varphi = \alpha(u) + \kappa(u)\ln u_x + \mu(u)\ln u_y. \tag{22}$$

Thus, the original classification problem (1) – (3) is reduced to three cases (20), (21) and (22). Now, we will analyze these three cases.

III. CASE 1. $\varphi = \alpha(\mathrm{u}) + \beta(\mathrm{u_x}) + \gamma(\mathrm{u_y})$

We substitute (20) in the equality (12). We get

$$\big(A(u,u_y) - u_y\alpha'(u)\big)\gamma'(u_y) = \big(B(u,u_x) - u_x\alpha'(u)\big)\beta'(u_x) = \mu(u).$$

Since u_x, u_y are independent, it follows that

$$A = \frac{\mu}{\gamma'} + u_y\alpha'(u), \quad B = \frac{\mu}{\beta'} + u_x\alpha'.$$

We replace A, B in (9), (10) and obtain

$$f = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}. \tag{23}$$

Substituting (23) in the equation (11) gives

$$\frac{\alpha'\mu}{\beta'\gamma'} - \mu^2\left(\frac{\gamma''}{\gamma'^2} + \frac{\beta''}{\beta'^2}\right)\frac{1}{\beta'\gamma'} + \alpha''u_xu_y + \mu'\left(\frac{u_x}{\gamma'} + \frac{u_y}{\beta'}\right) = F(\alpha+\beta+\gamma). \tag{24}$$

Thus, the problem (1)–(3) is reduced to (24).

By differentiating (24) with respect to u_x and u_y separately, we have

$$\begin{aligned} -\alpha'\mu\frac{\beta''}{\beta'^2\gamma'} - \mu^2\left(-\frac{\gamma''}{\gamma'^3}\frac{\beta''}{\beta'^2} + \frac{1}{\gamma'}\left(\frac{\beta''}{\beta'^3}\right)'\right) + \alpha''u_y + \mu'\left(\frac{1}{\gamma'} - u_y\frac{\beta''}{\beta'^2}\right) &= F'(\alpha+\beta+\gamma)\beta', \\ -\alpha'\mu\frac{\gamma''}{\beta'\gamma'^2} - \mu^2\left(\left(\frac{\gamma''}{\gamma'^3}\right)'\frac{1}{\beta'} - \frac{\beta''}{\beta'^3}\frac{\gamma''}{\gamma'^2}\right) + \alpha''u_x + \mu'\left(-u_x\frac{\gamma''}{\gamma'^2} + \frac{1}{\beta'}\right) &= F'(\alpha+\beta+\gamma)\gamma'. \end{aligned}$$

We exclude F' from these relations and obtain

$$-\alpha'\mu\frac{\beta''}{\beta'^2} - \mu^2\left(\frac{\beta''}{\beta'^3}\right)' + \alpha''u_y\gamma' - \mu'u_y\gamma'\frac{\beta''}{\beta'^2} = -\alpha'\mu\frac{\gamma''}{\gamma'^2} - \mu^2\left(\frac{\gamma''}{\gamma'^3}\right)' + \alpha''u_x\beta' - \mu'u_x\beta'\frac{\gamma''}{\gamma'^2}. \tag{25}$$

Applying $\dfrac{\partial^2}{\partial u_x \partial u_y}$ to both sides of (25), we get

$$\mu'\left((u_x\beta')'\left(\frac{\gamma''}{\gamma'^2}\right)' - (u_y\gamma')'\left(\frac{\beta''}{\beta'^2}\right)'\right) = 0. \tag{26}$$

It now follows that equation (26) is satisfied if one of the following conditions holds:

$$\mu'(u) = 0, \tag{27}$$

$$(u_x\beta')' = 0, \quad (u_y\gamma')' = 0, \tag{28}$$

$$(u_x\beta')' = 0, \quad \left(\frac{\beta''}{\beta'^2}\right)' = 0, \tag{29}$$

$$\left(\frac{\gamma''}{\gamma'^2}\right)' = 0, \quad (u_y\gamma')' = 0, \tag{30}$$

$$\left(\frac{\gamma''}{\gamma'^2}\right)' = 0, \quad \left(\frac{\beta''}{\beta'^2}\right)' = 0, \tag{31}$$

$$\frac{(u_x\beta')'}{\left(\frac{\beta''}{\beta'^2}\right)'} = \frac{(u_y\gamma')'}{\left(\frac{\gamma''}{\gamma'^2}\right)'} \neq 0. \tag{32}$$

Note that in the conditions (28) – (32), $\mu'(u) \neq 0$.

Now let us prove the following statement.

Lemma 1 *Under the condition* (27)*, we have that equations* (1)*,* (2) *and substitution* (3) *up to point transformations take the form:*

$$u_{xy} = 0, \quad v_{xy} = \exp v, \quad v = \alpha(u) + \ln(u_x u_y), \quad \alpha'' = \exp\alpha; \tag{33}$$

$$u_{xy} = u_x u_y, \quad v_{xy} = \exp v, \quad v = \alpha(u) + \ln(u_x u_y), \quad \alpha'' + \alpha' + 2 = \exp\alpha; \tag{34}$$

$$u_{xy} = -u_x u_y, \quad v_{xy} = 0, \quad v = \exp u + (a_1 + b_1)u + a_1 \ln u_x + b_1 \ln u_y; \tag{35}$$

$$\begin{gathered} u_{xy} = c\sqrt{u_x^2 + a_2}\sqrt{u_y^2 + b_2}, \quad v_{xy} = \frac{c^2}{2}\big(\exp v - a_2 b_2 \exp(-v)\big), \\ v = \ln\left(u_x + \sqrt{u_x^2 + a_2}\right)\left(u_y + \sqrt{u_y^2 + b_2}\right); \end{gathered} \tag{36}$$

$$u_{xy} = c\sqrt{1 - u_x^2}\sqrt{1 - u_y^2}, \quad v_{xy} = -c^2 \sin v, \quad v = \arcsin u_x + \arcsin u_y; \tag{37}$$

$$u_{xy} = c\sqrt{u_x u_y}, \quad v_{xy} = \frac{c^2 v}{4}, \quad v = \sqrt{u_x} + \sqrt{u_y}; \tag{38}$$

$$u_{xy} = c\sqrt{u_x}, \quad v_{xy} = \frac{c^2}{2}, \quad v = \sqrt{u_x} + u_y; \tag{39}$$

$$u_{xy} = c, \quad v_{xy} = 0, \quad v = u_x + u_y; \tag{40}$$

$$u_{xy} = c u_y\sqrt{1 - u_x^2}, \quad v_{xy} = -\mathrm{i}c^2 \exp v, \quad v = -\mathrm{i}\arcsin u_x + \ln u_y; \tag{41}$$

$$u_{xy} = \frac{a_1}{\gamma'(u_y)}, \quad v_{xy} = b_1, \quad v = u_x + \gamma(u_y) + u, \quad a_1 - a_1^2\frac{\gamma''}{\gamma'^2} = b_1\gamma'; \tag{42}$$

$$\begin{gathered} u_{xy} = a(u_x + c_7)(u_y + c_9), \quad v_{xy} = 0, \\ v = a_1 \ln(u_x + c_7) + b_1 \ln(u_y + c_9) + u, \quad a a_1 + a b_1 + 1 = 0; \end{gathered} \tag{43}$$

$$u_{xy} = 0, \quad v_{xy} = 0, \quad v = \beta(u_x) + \gamma(u_y) + u. \tag{44}$$

Here α'', α', 1 *are linearly independent functions,* c, c_1, c_2, c_7, c_9, $a_1 \neq 0$, $b_1 \neq 0$, $a \neq 0$, b_2, a_2 *are arbitrary constants.*

Proof. If the condition (27) holds, then $\mu(u) = c$, where c is a constant. We rewrite (25) and get

$$c\alpha' \frac{\beta''}{\beta'^2} + c^2 \left(\frac{\beta''}{\beta'^3}\right)' + \alpha'' u_x \beta' = c\alpha' \frac{\gamma''}{\gamma'^2} + c^2 \left(\frac{\gamma''}{\gamma'^3}\right)' + \alpha'' u_y \gamma' = \sigma(u). \tag{45}$$

From (45), we have

$$\sigma(u) = A_1 \alpha'(u) + B_1 \alpha''(u) + C_1, \tag{46}$$

where A_1, B_1, C_1 are constants. Rewriting (45) gives

$$\begin{aligned} \alpha' \left(c\tfrac{\beta''}{\beta'^2} - A_1\right) + \alpha'' \left(u_x \beta' - B_1\right) = C_1 - c^2 \left(\tfrac{\beta''}{\beta'^3}\right)', \\ \alpha' \left(c\tfrac{\gamma''}{\gamma'^2} - A_1\right) + \alpha'' \left(u_y \gamma' - B_1\right) = C_1 - c^2 \left(\tfrac{\gamma''}{\gamma'^3}\right)'. \end{aligned} \tag{47}$$

Suppose that 1, α', α'' are linearly independent functions, then

$$\begin{aligned} c\frac{\beta''}{\beta'^2} = A_1, \quad u_x\beta' = B_1, \quad C_1 - c^2 \left(\frac{\beta''}{\beta'^3}\right)' = 0, \\ c\frac{\gamma''}{\gamma'^2} = A_1, \quad u_y\gamma' = B_1, \quad C_1 - c^2 \left(\frac{\gamma''}{\gamma'^3}\right)' = 0. \end{aligned}$$

Therefore

$$\beta' = \frac{B_1}{u_x}, \quad \gamma' = \frac{B_1}{u_y}, \quad -\frac{c}{B_1} = A_1, \quad C_1 + \frac{c^2}{B_1^2} = 0.$$

We rewrite the equation (24) according to the above relations. We obtain

$$u_x u_y \left(\frac{c\alpha'}{B_1^2} + \frac{2c^2}{B_1^3} + \alpha''\right) = F(\alpha + \beta + \gamma). \tag{48}$$

Since 1, α', α'' are linearly independent functions, it follows that $F \neq 0$. By differentiating (48) with respect to u_x and using $\beta' = B_1/u_x$, we get

$$\frac{1}{u_x} = \frac{F'(\alpha + \beta + \gamma)}{F(\alpha + \beta + \gamma)} \cdot \frac{B_1}{u_x}.$$

Consequently

$$F(z) = C_1 \exp(z/B_1), \quad z = \alpha + \beta + \gamma. \tag{49}$$

We substitute (49) in (48) and obtain

$$\frac{c\alpha'}{B_1^2} + \frac{2c^2}{B_1^3} + \alpha'' = C_1 \exp\left(\frac{\alpha}{B_1} + \frac{C_2}{B_1} + \frac{C_3}{B_1}\right),$$

with $\beta = B_1 \ln u_x + C_2$, $\gamma = B_1 \ln u_y + C_3$. Thus, we have

$$
\begin{gathered}
u_{xy} = \frac{cu_xu_y}{B_1^2}, \quad v = \alpha(u) + B_1 \ln(u_xu_y) + C_2 + C_3, \quad v_{xy} = C_1 \exp(v/B_1), \\
c\frac{\alpha'}{B_1^2} + 2c^2\frac{1}{B_1^3} + \alpha'' = C_1 \exp\left(\frac{\alpha + C_2 + C_3}{B_1}\right).
\end{gathered}
\tag{50}
$$

After the replacement $(\alpha + C_2 + C_3)/B_1 \to \alpha$, $v/B_1 \to v$, equations (50) takes the form

$$
u_{xy} = \frac{cu_xu_y}{B_1^2}, \quad v = \alpha(u) + \ln(u_xu_y), \quad v_{xy} = \frac{C_1}{B_1}\exp v, \quad c\frac{\alpha'}{B_1^2} + 2c^2\frac{1}{B_1^4} + \alpha'' = \frac{C_1}{B_1}\exp\alpha. \tag{51}
$$

Finally, the replacements $v \to v + \ln(B_1/C_1)$, $\alpha \to \alpha + \ln(B_1/C_1)$, $c/B_1^2 \to c$ bring (51) to

$$
u_{xy} = cu_xu_y, \quad v = \alpha(u) + \ln(u_xu_y), \quad \alpha'' + c\alpha' + 2c^2 = \exp\alpha, \quad v_{xy} = \exp v. \tag{52}
$$

If $c = 0$, then equations (52) takes the form (33). If $c \neq 0$ then the replacements $cu \to u$, $\alpha \to \alpha + \ln c^2$ put (52) into (34). Thus, if 1, α', α'' are linearly independent functions then the case is reduced to (33) or (34).

Now, suppose that 1, α', α'' are linearly dependent functions. It means that

$$
C_1\alpha'' + C_2\alpha' + C_3 \cdot 1 = 0, \quad C_1^2 + C_2^2 + C_3^2 \neq 0.
$$

If $C_1 = 0$, then $C_2 \neq 0$ and we get

$$
\alpha' = c. \tag{53}
$$

If $C_1 \neq 0$, then

$$
\alpha'' = c_1\alpha' + c_2. \tag{54}
$$

Case (53) is a particular case of (54). Equation (54) has two families of solutions:

$$
\alpha = c_3u^2 + c_4u + c_5, \quad \alpha = \frac{1}{c_1}\exp c_1u + c_6u + c_7.
$$

The constants c_5, c_7 are reduced to zero by replacing $\beta + c_5 \to \beta$, $\beta + c_7 \to \beta$. Hence

$$
\alpha = c_2u^2 + c_3u \tag{55}
$$

or $\alpha = \frac{1}{c_1}\exp c_1u + c_4u$ which is equivalent to

$$
\alpha = \exp c_1u + c_4u, \quad c_1 \neq 0 \tag{56}
$$

by the replacement $u \to u + (\ln c_1)/c_1$ and reducing the constant to zero.

Now concentrate on the case (56) with taking into account that $\mu(u) = c$. Basic equation (24) becomes

$$\frac{c(c_1 \exp c_1 u + c_4)}{\beta'\gamma'} - c^2 \left(\frac{\gamma''}{\gamma'^2} + \frac{\beta''}{\beta'^2} \right) \frac{1}{\beta'\gamma'} + c_1^2 \exp(c_1 u) u_x u_y = F(\alpha + \beta + \gamma). \tag{57}$$

We differentiate (57) with respect to u and obtain

$$\frac{cc_1^2 \exp(c_1 u)}{\beta'\gamma'} + c_1^3 \exp(c_1 u) u_x u_y = F'(\alpha + \beta + \gamma)(c_1 \exp c_1 u + c_4).$$

Therefore

$$\frac{cc_1^2}{\beta'\gamma'} + c_1^3 u_x u_y = (c_1 + c_4 \exp(-c_1 u)) F'(\alpha + \beta + \gamma).$$

Applying $\dfrac{\partial}{\partial u}$ to both sides, we get

$$-c_1 c_4 \exp(-c_1 u) F'(\alpha + \beta + \gamma) + (c_1 + c_4 \exp(-c_1 u))(c_1 \exp c_1 u + c_4) F''(\alpha + \beta + \gamma) = 0.$$

Hence

$$(c_1 \exp c_1 u + c_4)^2 F''(\alpha + \beta + \gamma) = c_1 c_4 F'(\alpha + \beta + \gamma).$$

Consequently, we have two possibilities:

$$F'(\alpha + \beta + \gamma) = 0, \tag{58}$$

or

$$\frac{F''(\alpha + \beta + \gamma)}{F'(\alpha + \beta + \gamma)} = \frac{c_1 c_4}{(c_1 \exp c_1 u + c_4)^2}. \tag{59}$$

If (58) holds, then $F = c_5$, where c_5 is an arbitrary constant. In this case by using (57), we obtain

$$\frac{cc_1}{\beta'\gamma'} + c_1^2 u_x u_y = 0, \quad \frac{cc_4}{\beta'\gamma'} - c^2 \left(\frac{\gamma''}{\gamma'^2} + \frac{\beta''}{\beta'^2} \right) \frac{1}{\beta'\gamma'} = c_5.$$

By rewriting this relation, we get

$$\beta' = \frac{cc_1}{c_6 u_x}, \quad \gamma' = -\frac{c_6}{c_1^2 u_y}, \quad cc_4 - c^2 \left(\frac{c_1^2}{c_6} - \frac{c_6}{c_1 c} \right) = c_5 \beta'\gamma'.$$

These equations imply $c_5 = 0$ i.e. $F \equiv 0$. Thus, we have

$$\begin{gathered} u_{xy} = -c_1 u_x u_y, \quad v = \exp c_1 u + c_4 u + \frac{cc_1}{c_6} \ln u_x - \frac{c_6}{c_1^2} \ln u_y + c_7, \quad v_{xy} = 0, \\ c_4 - \frac{cc_1^2}{c_6} + \frac{c_6}{c_1} = 0, \quad c_1 \neq 0. \end{gathered} \tag{60}$$

Equations (60) by change of variables $c_1 u \to u$, $v \to v - cc_1 \ln(c_1)/c_6 + c_6 \ln(c_1)/c_1^2 + c_7$, $cc_1/c_6 \to a_1$, $-c_6/c_1^2 \to b_1$ are reduced to (35).

Now suppose that (59) is true. Differentiating (59) with respect to u_x gives

$$\left(\frac{F''}{F'}\right)'\beta' = 0.$$

Since $\beta' \neq 0$, we get

$$F(z) = c_6 \exp c_5 z + c_7, \quad c_5 c_6 \neq 0$$

or

$$F(z) = c_6 z + c_7, \quad c_6 \neq 0. \tag{61}$$

Change of variables $z \to z - (\ln c_6)/c_5$ implies

$$F(z) = \exp c_5 z + c_7, \quad c_5 \neq 0. \tag{62}$$

Let us consider the case (62). Under this case the equation (57) becomes

$$\frac{c(c_1 \exp c_1 u + c_4)}{\beta'\gamma'} - c^2\left(\frac{\gamma''}{\gamma'^2} + \frac{\beta''}{\beta'^2}\right)\frac{1}{\beta'\gamma'} + c_1^2 \exp(c_1 u) u_x u_y =$$
$$= \exp\big(c_5(\exp c_1 u + c_4 u)\big)\exp(c_5\beta)\exp(c_5\gamma).$$

This equality is not realized because $c_5 c_1 \neq 0$.

If (61) is true, then from (57) we obtain

$$\frac{c(c_1 \exp c_1 u + c_4)}{\beta'\gamma'} - c^2\left(\frac{\gamma''}{\gamma'^2} + \frac{\beta''}{\beta'^2}\right)\frac{1}{\beta'\gamma'} + c_1^2 \exp(c_1 u) u_x u_y = c_6(\exp c_1 u + c_4 u + \beta + \gamma) + c_7.$$

Differentiating this equation with respect u gives

$$\frac{c c_1^2 \exp c_1 u}{\beta'\gamma'} + c_1^3 u_x u_y \exp c_1 u = c_6(c_1 \exp c_1 u + c_4).$$

It implies

$$\frac{c c_1^2}{\beta'\gamma'} + c_1^3 u_x u_y = c_6 c_1, \quad c_6 c_4 = 0.$$

We see that this equality is realized if and only if $c_6 = 0$. Hence we get the case (35) considered above.

Assume that the condition (55) is satisfied. We substitute (55) in (24) with using $\mu(u) = c$. We obtain

$$\frac{c}{\beta'\gamma'}(2c_2 u + c_3) - c^2\left(\frac{\gamma''}{\gamma'^2} + \frac{\beta''}{\beta'^2}\right)\frac{1}{\beta'\gamma'} + 2c_2 u_x u_y = F(c_2 u^2 + c_3 u + \beta + \gamma).$$

Differentiating this equation with respect u and denoting $c_2u^2 + c_3 u + \beta + \gamma$ by z yields

$$2c_2 \frac{c}{\beta'\gamma'} = F'(z)(2c_2 u + c_3). \tag{63}$$

From the above equality we have the following cases:

$$c_2 = c_3 = 0, \tag{64}$$

$$c_2 \neq 0, \quad c_3 = 0, \tag{65}$$

$$c_2 = 0, \quad c_3 \neq 0, \tag{66}$$

$$c_2 c_3 \neq 0. \tag{67}$$

For the case (64) equations in (47) give

$$c_1 u_x - c^2 \frac{\beta''}{\beta'^3} = a_1, \quad c_1 u_y - c^2 \frac{\gamma''}{\gamma'^3} = b_1.$$

Here a_1, b_1 are arbitrary constants. Hence

$$\beta'(u_x) = \sqrt{-c^2} \frac{1}{\sqrt{c_1 u_x^2 - 2a_1 u_x + 2a_2}}, \tag{68}$$

$$\gamma'(u_y) = \sqrt{-c^2} \frac{1}{\sqrt{c_1 u_y^2 - 2b_1 u_y + 2b_2}}. \tag{69}$$

We integrate (68), (69) and obtain various formulae for determining the functions β, γ having the parameters c, c_1, a_i, b_i, $i = 1, 2$. We combine these formulae and get the following cases:

$$\begin{gathered} \beta(u_x) = a \ln\left(u_x - a_1 + \sqrt{(u_x - a_1)^2 + a_2}\right), \\ \gamma(u_y) = a \ln\left(u_y - b_1 + \sqrt{(u_y - b_1)^2 + b_2}\right); \end{gathered} \tag{70}$$

$$\beta(u_x) = a \arcsin \frac{u_x - a_1}{a_2}, \quad \gamma(u_y) = a \arcsin \frac{u_y - b_1}{b_2}; \tag{71}$$

$$\beta(u_x) = a \ln(u_x - a_1), \quad \gamma(u_y) = a \ln(u_y - b_1); \tag{72}$$

$$\beta(u_x) = a\sqrt{2a_2 - 2a_1 u_x}, \quad \gamma(u_y) = a\sqrt{2b_2 - 2b_1 u_x}; \tag{73}$$

$$\beta(u_x) = a\sqrt{2a_2 - 2a_1 u_x}, \quad \gamma(u_y) = a u_y; \tag{74}$$

$$\beta(u_x) = a u_x, \quad \gamma(u_y) = a u_y; \tag{75}$$

$$\beta(u_x) = \sqrt{\frac{c^2}{c_1}} \arcsin\left(\frac{u_x - \frac{a_1}{c_1}}{\sqrt{\frac{a_1^2 - 2a_2 c_1}{c_1^2}}}\right), \quad \gamma(u_y) = \sqrt{-\frac{c^2}{c_1}} \ln\left(u_y - \frac{b_1}{c_1}\right). \tag{76}$$

Here $a \neq 0$ is an arbitrary constant.

Substituting (70) in (23) and (24) with $\mu(u) = c$, $\alpha(u) = 0$ gives

$$\begin{gathered} u_{xy} = \frac{c}{a^2}\sqrt{(u_x - a_1)^2 + a_2}\sqrt{(u_y - b_1)^2 + b_2}, \quad v_{xy} = \frac{c^2}{2a^3}\exp(v/a) + b\exp(-v/a), \\ v = a \ln\left(u_x - a_1 + \sqrt{(u_x - a_1)^2 + a_2}\right)\left(u_y - b_1 + \sqrt{(u_y - b_1)^2 + b_2}\right), \end{gathered} \tag{77}$$

where $c^2a_2b_2/2a^3+b=0$. After the change of variables $u-a_1x-b_1y\to u$, $v/a\to v$, $c/a^2\to c$ equations (77) take the form (36).

If the condition (71) is true, then

$$u_{xy}=\frac{c}{a^2}\sqrt{a_2^2-(u_x-a_1)^2}\sqrt{b_2^2-(u_y-b_1)^2},\quad v_{xy}=\kappa\sin\frac{v}{a},\quad \kappa=-\frac{a_2b_2c^2}{a^3},$$
$$v=a\arcsin\frac{u_x-a_1}{a_2}+a\arcsin\frac{u_y-b_1}{b_2}.$$

These equations after the changes $v/a\to v$, $u-a_1x-b_1y\to u$ and then $a_2x\to x$, $b_2y\to y$, $c/a^2\to c$ take the form (37).

If (72) is true, then we get a particular case of (36).

Further, (73) gives

$$u_{xy}=\frac{c}{a^2a_1b_1}\sqrt{2a_2-2a_1u_x}\sqrt{2b_2-2b_1u_y},\quad v_{xy}=\frac{c^2}{a^4a_1b_1}v,$$
$$v=a\sqrt{2a_2-2a_1u_x}+a\sqrt{2b_2-2b_1u_y}.$$

After the transformaions $u-a_2x/a_1-b_2y/b_1\to -u$, $v/a\to v$, and then $x/2a_1\to x$, $y/2b_1\to y$,$4c/a^2\to -c$, these equations take the form (38).

By the expressions in (74) we have

$$u_{xy}=-\frac{c}{a_1a^2}\sqrt{2a_2-2a_1u_x},\quad v_{xy}=-\frac{c^2}{a_1a^3},\quad v=a\sqrt{2a_2-2a_1u_x}+au_y.$$

Changes $a_2x/a_1-u\to u$, then $v/a\to v$, $x/2a_1\to x$, $2c/a^2\to -c$, $-y\to y$ yield to equations (39).

Formula (75) determines $u_{xy}=c/a^2$, $v=au_x+au_y$, $v_{xy}=0$. These equations after changes $v/a\to v$, $c/a^2\to c$ take the form (40).

Substituting (76) in (23), (24) with $\mu(u)=c$, $\alpha(u)=0$ leads to

$$u_{xy}=-\frac{\mathrm{i}c_1}{c}\sqrt{-u_x^2+2\frac{a_1}{c_1}u_x-\frac{2a_2}{c_1}}\left(u_y-\frac{b_1}{c_1}\right),$$
$$v=\frac{c}{\sqrt{c_1}}\arcsin\left(\frac{u_x-\frac{a_1}{c_1}}{\sqrt{\frac{a_1^2-2a_2c_1}{c_1^2}}}\right)+\frac{\mathrm{i}c}{\sqrt{c_1}}\ln\left(u_y-\frac{b_1}{c_1}\right),$$
$$v_{xy}=-\frac{\sqrt{c_1}\sqrt{a_1^2-2a_2c_1}}{c}\exp\left(\frac{\sqrt{c_1}}{ic}v\right).$$

Changes $\sqrt{c_1}/(ic)v\to v$, $\sqrt{(a_1^2-2a_2c_1)/c_1^2}\to a_3$, $a_1/c_1\to a_3a$, $b_1/c_1\to b$, $a_3x\to x$ and, finally, $u-ax-by\to u$ give (41). Thus the case (64) is reduced to (36) – (41).

Now we consider the cases (65), (67). From the equality (63) we have

$$\ln\left(2c_2\frac{c}{\beta'\gamma'}\right) = \ln F'(z) + \ln(2c_2u + c_3).$$

We differentiate this equation with respect to u and get

$$0 = \frac{F''}{F'}(2c_2u + c_3) + \frac{2c_2}{2c_2u + c_3}. \tag{78}$$

Applying $\dfrac{\partial}{\partial u_x}$ to both sides of (78) gives $\left(\dfrac{F''}{F'}\right)' = 0$ i.e. $\dfrac{F''}{F'} = c_4$. Rewriting (78) yields to

$$c_4(2c_2u + c_3)^2 + 2c_2 = 0.$$

Hence $c_2 = 0$. This contradicts to the condition $c_2 \neq 0$.

If condition (66) is satisfied, then (63) gives $F(z) = c_4$. Here c_4 is an arbitrary constant. Rewriting (24) with using $\alpha = c_3u$, $\mu = c$ implies

$$c_3c - c^2\left(\frac{\gamma''}{\gamma'^2} + \frac{\beta''}{\beta'^2}\right) = c_4\beta'\gamma'. \tag{79}$$

We apply $\frac{\partial}{\partial u_x}$ to both sides of (79) and obtain

$$-c^2\left(\frac{\beta''}{\beta'^2}\right)' = c_4\beta''\gamma'. \tag{80}$$

Assume that $\beta'' = 0$ i.e. $\beta = c_5u_x + c_6$. Then (79) takes the form

$$c_3c - c^2\frac{\gamma''}{\gamma'^2} = c_4c_5\gamma'.$$

Thus, we have

$$u_{xy} = \frac{c}{c_5\gamma'(u_y)}, \quad v = c_5u_x + \gamma(u_y) + c_3u, \quad v_{xy} = c_4, \quad c_3c - c^2\frac{\gamma''}{\gamma'^2} = c_4c_5\gamma', \quad c_5 \neq 0.$$

The transformations $x/c_5 \to x$, $c_4c_5 \to c_2$, and then $v/c_3 \to v$, $\gamma/c_3 \to \gamma$, $c_3x \to x$, $c/c_3^2 \to a_1$, $c_2/c_3^2 \to b_1$ imply (42).

Suppose that $\beta'' \neq 0$, then

$$-c^2\frac{1}{\beta''(u_x)}\left(\frac{\beta''(u_x)}{\beta'^2(u_x)}\right)' = c_4\gamma'(u_y).$$

Since u_x, u_y are independent variables, we get

$$-c^2\frac{1}{\beta''}\left(\frac{\beta''}{\beta'^2}\right)' = c_5, \quad c_4\gamma' = c_5. \tag{81}$$

If $c_4 = 0$, then $c_5 = 0$ which yields to $c = 0$ or $\beta''/\beta'^2 = -c_6 \neq 0$. in the second case

$$\beta(u_x) = \frac{1}{c_6} \ln(c_6 u_x + c_7).$$

Inserting this formula in (79) and taking into account that $c_4 = 0$ gives the function $\gamma(u_y)$ as

$$\gamma(u_y) = \frac{1}{c_8} \ln(c_8 u_y + c_9).$$

Thus, we have equations

$$u_{xy} = c(c_6 u_x + c_7)(c_8 u_y + c_9), \quad v_{xy} = 0,$$
$$v = \frac{1}{c_6} \ln(c_6 u_x + c_7) + \frac{1}{c_8} \ln(c_8 u_y + c_9) + c_3 u, \quad cc_6 + cc_8 + c_3 = 0, \quad c_3 \neq 0.$$

After the transformations $v/c_3 \to v$, $x/c_6 \to x$, $y/c_8 \to y$, $1/(c_3c_6) \to a_1$, $1/(c_3c_8) \to b_1$, $cc_6c_8 \to a$, we obtain (43). Otherwise, if $c = 0$, $c_5 = c_4 = 0$ then we obtain (44).

Let us return to (81) and assume that $c_4 \neq 0$ i.e. we have

$$\gamma(u_y) = \frac{c_5}{c_4} u_y + c_6, \quad -c^2 \frac{\beta''}{\beta'^2} = c_5 \beta' + c_7.$$

We substitute the above relations in (79) and get $c_7 + cc_3 = 0$. Whence, we have

$$u_{xy} = \frac{cc_4}{c_5 \beta'(u_x)}, \quad v = \frac{c_5}{c_4} u_y + \beta(u_x) + c_3 u,$$
$$v_{xy} = c_4, \quad -c^2 \frac{\beta''}{\beta'^2} = c_5 \beta' + c_7, \quad c_7 + cc_3 = 0, \quad c_4 c_5 \neq 0.$$

It can be easily shown that this case is symmetrical case to (42) under $x \leftrightarrow y$.

Thus the case 1, α', α'' are linearly dependent gives equations (35) –(44). This completes the proof of Lemma.

Lemma 2 *Under the conditions* (28) *and* $\mu'(u) \neq 0$*, we have that equations* (1), (2) *and substitution* (3) *up to point transformations take the form:*

$$u_{xy} = \mu(u) u_x u_y, \quad v_{xy} = 0, \quad v = c_1 \ln u_x + c_2 \ln u_y + \alpha(u),$$
$$\mu'(c_1 + c_2) + \mu^2 (c_1 + c_2) + \alpha'' + \alpha' \mu = 0, \quad \mu' \neq 0; \tag{82}$$

$$u_{xy} = \mu(u) u_x u_y, \quad v_{xy} = \exp v, \quad v = \ln(u_x u_y) + \alpha(u),$$
$$2\mu' + 2\mu^2 + \alpha'' + \alpha' \mu = \exp \alpha; \tag{83}$$

Here $c_1 \neq 0$*,* $c_2 \neq 0$ *are arbitrary constants.*

Proof. If the condition (28) holds then the functions β, γ are equal to

$$\beta(u_x) = c_1 \ln u_x, \quad \gamma(u_y) = c_2 \ln u_y.$$

At the same time the equation (24) becomes

$$\begin{aligned} \mu'(u)u_x u_y \left(\frac{1}{c_2} + \frac{1}{c_1}\right) + \frac{\mu^2(u)u_x u_y}{c_1 c_2}\left(\frac{1}{c_2} + \frac{1}{c_1}\right) + \alpha''(u)u_x u_y + \alpha'(u)\mu(u)\frac{u_x u_y}{c_1 c_2} = \\ = F\big(c_1 \ln u_x + c_2 \ln u_y + \alpha(u)\big). \end{aligned} \tag{84}$$

Applying $\frac{\partial}{\partial u_x}$ to the both sides of (84) gives

$$\mu'(u)u_y \left(\frac{1}{c_2} + \frac{1}{c_1}\right) + \frac{\mu^2(u)u_y}{c_1 c_2}\left(\frac{1}{c_2} + \frac{1}{c_1}\right) + \alpha''(u)u_y + \alpha'(u)\mu(u)\frac{u_y}{c_1 c_2} = \frac{F' c_1}{u_x}.$$

Consequently, $F = c_1 F'$. We differentiating (84) with respect u_y in the same way. We get $F = c_2 F'$. This yields that $F' = 0$ or $c_2 = c_1$. If the first equation is true, then (84) turns out to be

$$u_x u_y \left(\mu'(u)\left(\frac{1}{c_2} + \frac{1}{c_1}\right) + \frac{\mu^2}{c_1 c_2}\left(\frac{1}{c_2} + \frac{1}{c_1}\right) + \alpha'' + \frac{\alpha'\mu}{c_1 c_2}\right) = c.$$

It implies that $c = 0$ and

$$u_{xy} = \frac{\mu(u)u_x u_y}{c_1 c_2}, \quad v = c_1 \ln u_x + c_2 \ln u_y + \alpha(u), \quad v_{xy} = 0,$$
$$\mu'\left(\frac{1}{c_2} + \frac{1}{c_1}\right) + \frac{\mu^2}{c_1 c_2}\left(\frac{1}{c_2} + \frac{1}{c_1}\right) + \alpha'' + \frac{\alpha'\mu}{c_1 c_2} = 0.$$

Finally, the transformation $\mu/c_1 c_2 \to \mu$ gives (82).

If $c_2 = c_1$ we have $F = c_3 \exp(v/c_1)$ then (84) becomes

$$\frac{2\mu' u_x u_y}{c_1} + \frac{2\mu^2 u_x u_y}{c_1^3} + \alpha'' u_x u_y + \frac{\alpha'\mu u_x u_y}{c_1^2} = c_3 u_x u_y \exp(\alpha(u)/c_1).$$

It follows that

$$u_{xy} = \frac{1}{c_1^2}\mu(u)u_x u_y, \quad v = c_1 \ln u_x u_y + \alpha(u), \quad v_{xy} = c_3 \exp(v/c_1),$$
$$\frac{2\mu'}{c_1} + \frac{2\mu^2}{c_1^3} + \alpha'' + \frac{\alpha'\mu}{c_1^2} = c_3 \exp(\alpha/c_1).$$

These equations after the changes $\mu/c_1^2 \to \mu$, $\alpha/c_1 \to \alpha$, $v/c_1 \to v$, then $v + \ln c \to v$, $\alpha + \ln c \to \alpha$, $c_3/c \to c_1$ takes the form (83).

Thus the condition (28) yields to equations (82), (83). This completes the proof of Lemma.

Lemma 3 *Assume that the condition* (30) *is satisfied but* (27), (28) *not. Then equations* (1), (2) *and substitution* (3) *up to point transformations take the form:*

$$u_{xy} = u, \quad v_{xy} = v, \quad v = c_1 u_y + c_2 u_x + c_3 u; \tag{85}$$

$$\begin{gathered} u_{xy} = \mu(u)(u_y + b)u_x, \quad v_{xy} = \exp v, \quad v = \ln(u_y + b) + \ln u_x + \alpha(u), \\ 2\mu' + 2\mu^2 + \alpha'' + \alpha'\mu = \exp\alpha, \quad 2\mu^2 + \mu' + \alpha'\mu = \exp\alpha; \end{gathered} \tag{86}$$

$$\begin{gathered} u_{xy} = \mu(u)(u_y + b)u_x, \quad v_{xy} = 0, \quad v = c_2\ln(u_y + b) + c_1\ln u_x + \alpha(u), \\ (\mu' + \mu^2)(c_1 + c_2) + \alpha'' + \alpha'\mu = 0, \quad c_1\mu' + \mu^2(c_1 + c_2) + \alpha'\mu = 0; \end{gathered} \tag{87}$$

$$\begin{gathered} u_{xy} = \mu(u)u_x, \quad v_{xy} = 0, \quad v = u_y - \ln u_x + \alpha(u), \\ \alpha'' + \mu' = 0, \quad \mu^2 - \mu' + \alpha'\mu = 0. \end{gathered} \tag{88}$$

Here $c_1 c_2 b \neq 0$, c_3 *are arbitrary constants.*

Proof. If (30) holds, then we have the following three possibilities for the functions β, γ:

$$\gamma(u_y) = c_1 u_y + c_2, \quad \beta(u_x) = c_3 u_x + c_4, \tag{89}$$

$$\gamma(u_y) = -\frac{1}{c_1}\ln(a_1 u_y + b_1), \quad \beta(u_x) = -\frac{1}{c_2}\ln(a_2 u_x + b_2), \tag{90}$$

$$\gamma(u_y) = c_1 u_y + c_2, \quad \beta(u_x) = -\frac{1}{c_3}\ln(a u_x + b). \tag{91}$$

With (89) the equation (24) takes the form

$$\frac{\mu'(u)u_y}{c_3} + \frac{\mu'(u)u_x}{c_1} + \alpha''(u)u_x u_y + \frac{\alpha'(u)\mu(u)}{c_1 c_3} = F(c_1 u_y + c_3 u_x + \alpha(u)). \tag{92}$$

By applying $\frac{\partial}{\partial u_x}$, $\frac{\partial}{\partial u_y}$ to both sides of (92), we get

$$\frac{\mu'}{c_1} + \alpha'' u_y = F' c_3, \quad \frac{\mu'}{c_3} + \alpha'' u_x = F' c_1. \tag{93}$$

Excluding F' gives

$$\alpha''(c_1 u_y - c_3 u_x) = 0.$$

Hence $\alpha'' = 0$ are $\alpha = c_2 u + c_4$. Furthermore, by using (93), we get $F' = \mu'/c_1 c_3$. We differentiate (92) with respect to u

$$\frac{\mu'' u_y}{c_3} + \frac{\mu'' u_x}{c_1} + \frac{c_2\mu'}{c_1 c_3} = F' c_2.$$

When we insert $F' = \mu'/c_1 c_3$ in this equation, we obtain $\mu'' = 0$ i.e. $\mu(u) = c_5 u + c_6$. Moreover,

$$F(z) = \frac{c_5}{c_1 c_3} z + c_7.$$

If we use the above relations in (92), we get

$$\frac{c_2c_6}{c_1c_3}=\frac{c_5c_4}{c_1c_3}+c_7.$$

Thus, we have

$$\begin{gathered} u_{xy}=\frac{c_5u+c_6}{c_1c_3}, \quad v_{xy}=\frac{c_5}{c_1c_3}v+c_7, \\ v=c_1u_y+c_3u_x+c_2u+c_4, \quad \frac{c_2c_6}{c_1c_3}=\frac{c_5c_4}{c_1c_3}+c_7. \end{gathered}$$

The transformations $u+c_6/c_5\to cu$, $v+c_7\frac{c_1c_3}{c_5}\to cv$, $c_5/c_1\to c_3$ give equations (85).

By (90) the equation (24) takes the form

$$\begin{aligned} -c_2\mu' u_y\frac{a_2u_x+b_2}{a_2}-\mu^2\frac{c_1c_2^2}{a_1a_2}(a_2u_x+b_2)(a_1u_y+b_1)-\mu^2\frac{c_1^2c_2}{a_1a_2}(a_1u_y+b_1)(a_2u_x+b_2)- \\ -c_1\mu' u_x\frac{a_1u_y+b_1}{a_1}+\alpha'' u_xu_y+\alpha'\mu\frac{c_1c_2}{a_1a_2}(a_2u_x+b_2)(a_1u_y+b_1)= \\ =F\left(-\frac{1}{c_1}\ln(a_1u_y+b_1)-\frac{1}{c_2}\ln(a_2u_x+b_2)+\alpha(u)\right). \end{aligned} \tag{94}$$

Applying $\frac{\partial}{\partial u_x}$ to both sides of (94) gives

$$\begin{gathered} -c_2\mu' u_y-\mu^2\frac{c_1c_2^2}{a_1}(a_1u_y+b_1)-c_1\mu'\frac{a_1u_y+b_1}{a_1}-\mu^2\frac{c_1^2c_2}{a_1}(a_1u_y+b_1)+ \\ +\alpha'' u_y+\alpha'\mu\frac{c_1c_2}{a_1}(a_1u_y+b_1)=F'\left(-\frac{1}{c_2}\right)\frac{a_2}{a_2u_x+b_2}. \end{gathered}$$

It is clear that

$$F'\left(-\frac{1}{c_2}\right)-F=-c_1\mu'\frac{a_1u_y+b_1}{a_1}\frac{b_2}{a_2}+\alpha'' u_y\frac{b_2}{a_2}. \tag{95}$$

In the same way, by differentiating (94) with respect u_y, we obtain

$$F'\left(-\frac{1}{c_1}\right)-F=-c_2\mu'\frac{a_2u_x+b_2}{a_2}\frac{b_1}{a_1}+\alpha'' u_x\frac{b_1}{a_1}. \tag{96}$$

We differentiate (95) by u_x and (96) by u_y. We get

$$F''\left(-\frac{1}{c_2}-F'\right)=0, \quad F''\left(-\frac{1}{c_1}-F'\right)=0.$$

These equations yield to $F''(c_2-c_1)=0$. Thus, we have two possibilities:

$$F''=0 \tag{97}$$

or

$$c_2=c_1. \tag{98}$$

If $c_1 = c_2 = c$, then

$$F(z) = -\frac{1}{c}\exp(-cz + c_7) + c_8. \tag{99}$$

We substitute (99) in the equation (94). We get

$$\begin{gathered}
-\mu' u_y c\left(u_x + \frac{b_2}{a_2}\right) - 2\mu^2\frac{c^3}{a_1a_2}(a_2u_x + b_2)(a_1u_y + b_1) - \mu' u_x c\left(u_y + \frac{b_1}{a_1}\right) + \alpha'' u_x u_y + \\
+\alpha'\mu\frac{c^2}{a_1a_2}(a_2u_x + b_2)(a_1u_y + b_1) = -\frac{1}{c}(a_2u_x + b_2)(a_1u_y + b_1)\exp(-c\alpha + c_7) + c_8.
\end{gathered}$$

Since u, u_x, u_y are independent variables, we obtain

$$\begin{aligned}
-2c\mu' - 2\mu^2c^3 + \alpha'' + \alpha'\mu c^2 &= -\frac{a_1a_2}{c}\exp(-c\alpha + c_7),\\
-2\mu^2\frac{c^3}{a_1}b_1 - \mu' c\frac{b_1}{a_1} + \alpha'\mu c^2\frac{b_1}{a_1} &= -\frac{1}{c}a_2b_1\exp(-c\alpha + c_7),\\
-\mu' c\frac{b_2}{a_2} - 2\mu^2c^3\frac{b_2}{a_2} + \alpha'\mu c^2\frac{b_2}{a_2} &= -\frac{1}{c}a_1b_2\exp(-c\alpha + c_7),\\
-2\mu^2c^3\frac{b_1b_2}{a_1a_2} + \alpha'\mu c^2\frac{b_1b_2}{a_1a_2} &= -\frac{1}{c}b_1b_2\exp(-c\alpha + c_7) + c_8.
\end{aligned} \tag{100}$$

Note that $b_1^2 + b_2^2 \neq 0$, otherwise we have the condition (28) holds, which is a contradiction to the assumption of the Lemma. If $b_2 = 0$, $b_1 \neq 0$, then $c_8 = 0$ and

$$\begin{gathered}
u_{xy} = \frac{\mu(u)c^2}{a_1}(a_1u_y + b_1)u_x, \quad v_{xy} = -\frac{1}{c}\exp(-cv + c_7),\\
v = -\frac{1}{c}\ln(a_1u_y + b_1) - \frac{1}{c}\ln(a_2u_x) + \alpha(u),\\
-2c\mu' - 2\mu^2c^3 + \alpha'' + \alpha'\mu c^2 = -\frac{a_1a_2}{c}\exp(-c\alpha + c_7),\\
-2\mu^2c^3 - \mu' c + \alpha'\mu c^2 = -\frac{a_1a_2}{c}\exp(-c\alpha + c_7).
\end{gathered}$$

The transformations $-cv + c_7 \to v$, $-c\alpha + c_7 + \ln(a_1a_2) \to \alpha$, $\mu c^2 \to \mu$, $b_1/a_1 \to b$ give equations (86). If $b_1b_2 \neq 0$, then it is not hard to prove that (100) is inconsistent.

Assume that (97) is satisfied, then

$$F(z) = cz + p.$$

Inserting this expression in (94) leads to

$$\begin{gathered}
-c_2\mu' u_y\left(u_x + \frac{b_2}{a_2}\right) - \mu^2c_1c_2^2\left(u_x + \frac{b_2}{a_2}\right)\left(u_y + \frac{b_1}{a_1}\right) - c_1\mu' u_x\left(u_y + \frac{b_1}{a_1}\right) - \\
-\mu^2c_1^2c_2\left(u_x + \frac{b_2}{a_2}\right)\left(u_y + \frac{b_1}{a_1}\right) + \alpha'' u_x u_y + \alpha'\mu c_1c_2\left(u_x + \frac{b_2}{a_2}\right)\left(u_y + \frac{b_1}{a_1}\right) = \\
= c\left(-\frac{1}{c_1}\ln(a_1u_y + b_1) - \frac{1}{c_2}\ln(a_2u_x + b_2) + \alpha(u)\right) + p.
\end{gathered}$$

Therefore $c = 0$. Since u, u_x, u_y are independent variables, we obtain

$$\begin{gathered}
-c_2\mu' - \mu^2 c_1 c_2^2 - c_1\mu' - \mu^2 c_1^2 c_2 + \alpha'' + \alpha'\mu c_1 c_2 = 0, \\
-\mu^2 c_1 c_2^2 \frac{b_1}{a_1} - c_1\mu' \frac{b_1}{a_1} - \mu^2 c_1^2 c_2 \frac{b_1}{a_1} + \alpha'\mu c_1 c_2 \frac{b_1}{a_1} = 0, \\
-c_2\mu' \frac{b_2}{a_2} - \mu^2 c_1 c_2^2 \frac{b_2}{a_1} - \mu^2 c_1^2 c_2 \frac{b_2}{a_1} + \alpha'\mu c_1 c_2 \frac{b_2}{a_2} = 0, \\
-\mu^2 c_1 c_2^2 - \mu^2 c_1^2 c_2 + \alpha'\mu c_1 c_2 \frac{b_1 b_2}{a_1 a_2} = p.
\end{gathered}$$

Note that $b_1^2 + b_2^2 \neq 0$, in the converse case we have (28) satisfied which is a contradiction. If $b_2 = 0$, $b_1 \neq 0$, then $p = 0$ and

$$\begin{gathered}
u_{xy} = \mu(u)\frac{c_1 c_2}{a_1}(a_1 u_y + b_1)u_x, \quad v_{xy} = 0, \\
v = -\frac{1}{c_1}\ln(a_1 u_y + b_1) - \frac{1}{c_2}\ln(a_2 u_x) + \alpha(u), \\
\mu'(c_1 + c_2) + \mu^2 c_1 c_2 (c_1 + c_2) - \alpha'' - \alpha'\mu c_1 c_2 = 0, \\
c_1\mu' + \mu^2 c_1 c_2 (c_1 + c_2) - \alpha'\mu c_1 c_2 = 0.
\end{gathered}$$

The replacements $c_1 c_2 \mu \to \mu$, $-c_1 c_2 \alpha + c_2 \ln a_1 + c_1 \ln a_2 \to \alpha$, $-c_1 c_2 v \to v$, $b_1/a_1 \to b$ give equations (87). If $b_1 b_2 \neq 0$, then we get inconsistent system for α, μ.

Suppose that the condition (91) is true. We rewrite (24) according to that and get

$$\begin{gathered}
-\frac{c_2}{a}\mu' u_y (a u_x + b) + \frac{c_2^2}{a c_1}(a u_x + b) + \frac{1}{c_1}\mu' u_x + \alpha'' u_x u_y + \\
+\alpha'\mu(a u_x + b)\left(-\frac{c_2}{c_1 a}\right) = F\left(c_1 u_y - \frac{1}{c_2}\ln(a u_x + b) + \alpha(u)\right).
\end{gathered} \tag{101}$$

Applying $\frac{\partial}{\partial u_x}$, $\frac{\partial}{\partial u_y}$ to both sides of (101) gives

$$-c_2\mu' u_y + \frac{c_2^2}{c_1}\mu^2 + \frac{1}{c_1}\mu' + \alpha'' u_y - \frac{c_2}{c_1}\alpha'\mu = F'\left(-\frac{1}{c_2}\right)\frac{a}{a u_x + b}, \tag{102}$$

$$-\frac{c_2}{a}\mu'(a u_x + b) + \alpha'' u_x = F' c_1. \tag{103}$$

If $F' = 0$, then $F = c_3$ and

$$-c_2\mu' + \alpha'' = 0, \quad c_2^2\mu^2 + \mu' - c_2\alpha'\mu = 0, \quad \mu' b = 0.$$

Substituting these equations in (101), we get that $c_3 = 0$. Thus, we have

$$\begin{gathered}
u_{xy} = -\frac{c_2}{c_1}\mu(u)u_x, \quad v_{xy} = 0, \quad v = c_1 u_y - \frac{1}{c_2}\ln a u_x + \alpha(u), \\
\alpha'' = c_2\mu', \quad c_2^2\mu^2 + \mu' - c_2\alpha'\mu = 0.
\end{gathered}$$

By a suitable point replacement we get equations (88).

Assume that $F' \neq 0$. Excluding F' from (102), (103), we obtain

$$c_2^2\left(\frac{au_x+b}{a}\right)\mu' u_y-\frac{c_2^3}{c_1}\left(\frac{au_x+b}{a}\right)\mu^2-\frac{c_2}{c_1}\left(\frac{au_x+b}{a}\right)\mu'-c_2\left(\frac{au_x+b}{a}\right)\alpha'' u_y+$$
$$+\frac{c_2^2}{c_1}\alpha'\mu\left(\frac{au_x+b}{a}\right)=-\frac{c_2}{c_1 a}(au_x+b)\mu'+\frac{\alpha''}{c_1}u_x.$$

Since u, u_x, u_y are independent variables, we get

$$\begin{gathered}
c_2^2\mu'-c_2\alpha''=0,\\
-\frac{c_2^3}{c_1}\mu^2-\frac{c_2}{c_1}\mu'+\frac{c_2^2}{c_1}\alpha'\mu=-\frac{c_2}{c_1}\mu'+\frac{\alpha''}{c_1},\\
\frac{c_2^2 b}{a}\mu'-c_2\frac{b}{a}\alpha''=0,\\
-\frac{c_2^3}{c_1}\frac{b}{a}\mu^2+\frac{c_2^2}{c_1}\alpha'\mu\frac{b}{a}=0.
\end{gathered}\tag{104}$$

If $b=0$, then the equation (101) takes the form

$$-c_2\mu' u_x u_y+\frac{c_2^2}{c_1}\mu^2 u_x+\frac{1}{c_1}\mu' u_x+\alpha'' u_x u_y-\frac{c_2}{c_1}\alpha'\mu u_x=F\left(c_1 u_y-\frac{1}{c_2}\ln(au_x)+\alpha(u)\right).$$

Differentiating this equation with respect to u_x, we obtain

$$\frac{c_2^2}{c_1}\mu^2-c_2\mu' u_y+\frac{1}{c_1}\mu'+\alpha'' u_y-\frac{c_2}{c_1}\alpha'\mu=-\frac{1}{c_2}F'\frac{1}{u_x}.$$

Consequently, $F+F'/c_2=0$ or $F(z)=c_3\exp(-c_2 z)$. Therefore

$$-c_2\mu' u_x u_y+\frac{c_2^2}{c_1}\mu^2 u_x+\frac{1}{c_1}\mu' u_x+\alpha'' u_x u_y-\frac{c_2}{c_1}\alpha'\mu u_x=c_3\exp(-c_2 c_1 u_y)au_x\exp(\alpha).$$

Since $c_3\neq 0$, $a\neq 0$, this equality is not realized. In the converse case $b\neq 0$, the equation (104) gives

$$c_2\mu'-\alpha''=0,\quad -c_2^3\mu^2+c_2^2\alpha'\mu=\alpha'',\quad -c_2\mu^2+\alpha'\mu=0.$$

We subtract first equation from second one and consider third equation. We get that $\mu'=0$. This contradicts to the hypothesis.

Thus, the condition (30) yields to equations (85)– (88). This completes the proof.

Lemma 4 *Suppose that the condition* (29) *holds but* (27), (28), (30) *not. Then equations*

(1), (2) *and substitution* (3) *up to point transformations take the form:*

$$\begin{gathered}u_{xy}=\frac{\mu(u)u_x}{\gamma'(u_y)},\quad v_{xy}=0,\quad v=\ln u_x+\gamma(u_y)+\alpha(u),\\ c_3+\frac{\gamma''}{\gamma'^2}+c_4\gamma' u_y=0,\quad \alpha''+\mu'+c_4\mu^2=0,\quad c_3\mu^2+\mu'+\mu^2+\alpha'\mu=0;\end{gathered}\tag{105}$$

$$\begin{gathered}u_{xy}=\frac{u_x}{(au+b)\gamma'(u_y)},\quad v_{xy}=\exp v,\quad v=\ln u_x+\gamma(u_y)-2\ln(au+b)+\ln(-c_4),\\ c_1+\frac{\gamma''}{\gamma'^2}+c_3\gamma' u_y=c_4\gamma'\exp\gamma,\quad c_1+1-3a=0,\quad c_3+2a^2-a=0.\end{gathered}\tag{106}$$

Proof. In the case of (29) we have $\beta=c_1\ln u_x+c_2$. After replacement $c_2+\alpha\to\alpha$ we get

$$\beta=c_1\ln u_x.\tag{107}$$

Substituting (107) in (24)

$$\frac{\alpha'\mu u_x}{c_1\gamma'}-\frac{\mu^2u_x}{c_1\gamma'}\left(\frac{\gamma''}{\gamma'^2}-\frac{1}{c_1}\right)+\alpha'' u_xu_y+\mu'\left(\frac{u_x}{\gamma'}+\frac{u_xu_y}{c_1}\right)=F(\alpha+\beta+\gamma).\tag{108}$$

Applying $\frac{\partial}{\partial u_x}$ to both sides of (108) gives

$$\frac{\alpha'\mu}{c_1\gamma'}-\frac{\mu^2}{c_1\gamma'}\left(\frac{\gamma''}{\gamma'^2}-\frac{1}{c_1}\right)+\alpha'' u_y+\mu'\left(\frac{1}{\gamma'}+\frac{u_y}{c_1}\right)=F'\left(\frac{c_1}{u_x}\right).\tag{109}$$

Note that (108), (109) yield to $F=F'\dfrac{c_1}{u_x}$ or

$$F(z)=c_2\exp(z/c_1).\tag{110}$$

By substituting (110) in (108), we have

$$u_x\left(\frac{\alpha'\mu}{c_1\gamma'}-\frac{\mu^2}{c_1\gamma'}\left(\frac{\gamma''}{\gamma'^2}-\frac{1}{c_1}\right)+\alpha'' u_y+\mu'\left(\frac{1}{\gamma'}+\frac{u_y}{c_1}\right)\right)=c_2u_x\exp(\gamma/c_1)\exp(\alpha/c_1).$$

Hence

$$\mu' c_1+\alpha'\mu+\frac{\mu^2}{c_1}-\mu^2\frac{\gamma''}{\gamma'^2}+(\alpha'' c_1+\mu')\gamma' u_y=c_2c_1\gamma'\exp(\gamma/c_1)\exp(\alpha/c_1).$$

Let us fix u, then γ is determined from

$$c_3+\frac{\gamma''}{\gamma'^2}+c_4\gamma' u_y=c_1c_5\gamma'\exp(\gamma/c_1).$$

Moreover

$$\alpha'\mu+\frac{\mu^2}{c_1}+c_1\mu'+c_3\mu^2+\gamma u_y(c_1\alpha''+\mu'+c_4\mu^2)-c_1\gamma'\exp(\gamma/c_1)(c_5\mu^2+c_2\exp(\alpha/c_1))=0.$$

If $u_y = \kappa \exp(\gamma/c_1)$, then $\gamma = c_1 \ln(u_y/\kappa)$ and $(\gamma' u_y)' = 0$. Since this equation contradicts to the condition of the Lemma, we get that u_y and $\exp(\gamma/c_1)$ are linearly independent and

$$c_1\alpha'' + \mu' + c_4\mu^2 = 0, \quad c_5\mu^2 + c_2 \exp(\alpha/c_1) = 0, \quad c_3\mu^2 + c_1\mu' + \frac{\mu^2}{c_1} + \alpha'\mu = 0.$$

Clearly, the constants $c_5 = 0$, so $c_2 = 0$ and

$$u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v = \beta(u_x) + \gamma(u_y) + \alpha(u), \quad v_{xy} = 0,$$
$$\beta' = \frac{c_1}{u_x}, \quad c_3 + \frac{\gamma''}{\gamma'^2} + c_4\gamma' u_y = 0, \quad c_1\alpha'' + \mu' + c_4\mu^2 = 0, \quad c_3\mu^2 + c_1\mu' + \frac{\mu^2}{c_1} + \alpha'\mu = 0.$$

The replacements $v/c_1 \to v$, $\alpha/c_1 \to \alpha$, $\gamma/c_1 \to \gamma$, $\mu/c_1^2 \to \mu$, $c_4c_1^2 \to c_4$, $c_1c_3 \to c_3$ give equations (105).

If $c_5 \neq 0$, then we have

$$u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v = \beta(u_x) + \gamma(u_y) + \alpha(u), \quad v_{xy} = c_2 \exp(v/c_1),$$

with

$$\beta' = \frac{c_1}{u_x}, \quad c_3 + \frac{\gamma''}{\gamma'^2} + c_4\gamma' u_y = c_1c_5\gamma' \exp(\gamma/c_1),$$
$$\alpha = 2c_1 \ln(-2c_1) - 2c_1 \ln\left(-\frac{2}{3}\sqrt{-\frac{c_2}{c_5}}\frac{c_3c_1+1}{c_1}u + c_6\right),$$
$$\mu = \sqrt{-\frac{c_2}{c_5}}\left(\frac{-2c_1}{-\frac{2}{3}\sqrt{-\frac{c_2}{c_5}}\left(\frac{c_3c_1+1}{c_1}\right)u + c_6}\right),$$
$$\frac{2}{9}\left(\frac{c_3c_1+1}{c_1}\right)^2 - \frac{1}{3}\left(\frac{c_3c_1+1}{c_1^2}\right) + c_4 = 0.$$

We make the substitutions with proper transformations and get equations (106).

Thus under the condition (29) we have equations (105), (106). This completes the proof.

Lemma 5 *Suppose that the condition* (32) *holds but* (27)–(31) *not, then equations* (1), (2) *and* (3) *up to point transformations take the form:*

$$\begin{gathered}u_{xy} = -\frac{1}{u\beta'(u_x)\gamma'(u_y)}, \quad v_{xy} = 0, \quad v = \beta(u_x) + \gamma(u_y),\\ \frac{\beta''}{\beta'^2} = u_x\beta' + c_1, \quad \gamma''/\gamma'^2 = u_y\gamma' - c_1;\end{gathered} \tag{111}$$

$$\begin{gathered}u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v_{xy} = \exp v, \quad v = \beta(u_x) + \gamma(u_y) + \alpha(u),\\ u_x + \frac{1}{\beta'(u_x)} = \exp(\beta), \quad u_y + \frac{1}{\gamma'(u_y)} = \exp\gamma, \quad \alpha'' = \exp\alpha, \quad \mu = (\exp\alpha)/\alpha';\end{gathered} \tag{112}$$

$$\begin{gathered}u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v_{xy} = \exp v, \quad v = \beta(u_x) + \gamma(u_y) + \alpha(u),\\ -cu_x + \frac{1}{\beta'(u_x)} = \exp\beta, \quad -cu_y + \frac{1}{\gamma'(u_y)} = \exp\gamma,\\ \alpha'\mu + 2\mu^2(c+1) = \exp\alpha, \quad \alpha'^2 = 2c^2\exp\alpha, \quad c = -\tfrac{1}{2}, -2;\end{gathered} \tag{113}$$

$$\begin{gathered}u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v = \beta(u_x) + \gamma(u_y) + \alpha(u), v_{xy} = \exp v + \exp(-v),\\ A_1\exp\beta + B_1\exp(-\beta) = u_x, \quad A_2\exp\gamma + B_2\exp(-\gamma) = u_y,\\ \alpha'' = \frac{1}{4}\left(\frac{\exp(-\alpha)}{B_1B_2} + \frac{\exp\alpha}{A_1A_2}\right), \quad \mu = \frac{\alpha''}{\alpha'};\end{gathered} \tag{114}$$

$$\begin{gathered}u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v = \beta(u_x) + \gamma(u_y) + \alpha(u), \quad v_{xy} = \exp v + \exp(\sigma v),\\ A_1\exp\beta + B_1\exp(\sigma\beta) = u_x, \quad A_2\exp\gamma + B_2\exp(\sigma\gamma) = u_y,\\ \alpha'^2 = \frac{2}{(\sigma-1)^2}\left(\sigma^2\frac{\exp\alpha}{A_1A_2} + \frac{1}{\sigma}\frac{\exp(\sigma\alpha)}{B_1B_2}\right),\\ 2(\sigma+1)\mu^2 + \alpha'\mu - \frac{1}{(\sigma-1)^2}\left(\frac{\exp\alpha}{A_1A_2} + \frac{\exp(\sigma\alpha)}{B_1B_2}\right) = 0, \quad \sigma = -\frac{1}{2}, -2.\end{gathered} \tag{115}$$

Proof. Since u_x, u_y are independent variables, (32) yields to

$$\frac{(u_x\beta')'}{\left(\frac{\beta''}{\beta'^2}\right)'} = c, \quad \frac{(u_y\gamma')'}{\left(\frac{\gamma''}{\gamma'^2}\right)'} = c, \quad c \neq 0.$$

Integrating these equations, we obtain

$$\frac{\beta''}{\beta'^2} = cu_x\beta' + c_1, \quad \frac{\gamma''}{\gamma'^2} = cu_y\gamma' + c_2. \tag{116}$$

By rewriting (24) and using (116), we get

$$\begin{gathered}\frac{1}{\beta'}\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(c_1 + cu_y\gamma' + c_2) + \mu' u_y\right) +\\ +u_x\left(-\frac{c\mu^2}{\gamma'} + \alpha'' u_y + \frac{\mu'}{\gamma'}\right) = F(\alpha+\beta+\gamma).\end{gathered} \tag{117}$$

Let us fix u_x and u_y, then $F(\beta + c_3) = c_4 u_x + c_5/\beta'$. After transformation $\beta + c_3 \to \beta$ we get

$$F(\beta) = c_4 u_x + \frac{c_5}{\beta'}. \tag{118}$$

Applying $\frac{\partial}{\partial u_x}$ to both sides of (118) and using (116) give rise to

$$F'(\beta) = -cc_5 u_x + \frac{c_4 - c_1 c_5}{\beta'}.$$

We differentiate this equation with respect to u_x and we get

$$F''(\beta) = -c(c_4 - c_1 c_5) u_x - \frac{cc_5 + c_1(c_4 - c_1 c_5)}{\beta'}.$$

Now, we see that

$$F'' = c_7 F' + c_8 F. \tag{119}$$

General solution of (119) has the forms

$$F(v) = A_1 \exp(\sigma_1 v) + B_1 \exp(\sigma_2 v), \quad \sigma_1 \neq \sigma_2$$

or

$$F(v) = (A_2 + B_2 v) \exp(\sigma v).$$

Considering different values of the constants A_i, B_i, $i = 1, 2$, we obtain different formulae for F:

$$F(v) = 0, \tag{120}$$

$$F(v) = 1, \tag{121}$$

$$F(v) = v, \tag{122}$$

$$F(v) = v \exp v, \tag{123}$$

$$F(v) = \exp v, \tag{124}$$

$$F(v) = \exp v + 1, \tag{125}$$

$$F(v) = \exp v + \exp(\sigma v). \tag{126}$$

By fixing different values u, u_y in (117), we obtain a set of equations

$$\alpha_i u_x + \frac{\beta_i}{\beta'(u_x)} = F\left(\beta(u_x) + \gamma_i\right). \tag{127}$$

Here α_i, β_i, γ_i are constants.

(i) Vectors $(\alpha_{\mathbf{i}}, \beta_{\mathbf{i}})$ are proportional. This means that there exists a set of numbers μ_i, such that

$$(\alpha_i, \beta_i) = \mu_i(\alpha_1, \beta_1), \quad \mu_x = 1.$$

Let us rewrite (127),

$$\mu_i \left(\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)} \right) = F(\beta + \gamma_i). \tag{128}$$

(i.1) If (120) holds then the equation (128) takes the form

$$\mu_i \left(\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)} \right) = 0. \tag{129}$$

Suppose that $\alpha_1 = \beta_1 = 0$. Using (117), we get

$$\mu' - c\mu^2 + \alpha'' u_y \gamma' = 0, \quad \alpha' \mu - \mu^2(c_1 + c_2 + c u_y \gamma') + \mu' u_y \gamma' = 0. \tag{130}$$

If $\alpha'' = 0$ i.e. $\alpha = \epsilon u + \delta$, the first equation of (130) gives

$$\mu(u) = -\frac{1}{cu + \kappa}.$$

Substituting the function μ into the second equation of (130) gives

$$\frac{\epsilon}{cu + \kappa} + \frac{c_1 + c_2}{(cu + \kappa)^2} = 0.$$

Therefore $\epsilon = 0$, $c_2 = -c_1$ and

$$u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v = \beta(u_x) + \gamma(u_y) + \alpha(u), \quad v_{xy} = 0,$$
$$\mu(u) = -\frac{1}{cu + \kappa}, \quad \alpha(u) = \delta, \quad \frac{\beta''}{\beta'^2} = c u_x \beta' + c_1, \quad \frac{\gamma''}{\gamma'^2} = c u_y \gamma' - c_1.$$

The transformations $\beta \to a\beta$, $\gamma \to a\gamma$, $a^2 c \to 1$, $u + \kappa/c \to u$, $v - \delta \to av$, $ac_1 \to c_1$ give equations (111).

Now assume that $\alpha'' \neq 0$. The first equation of (130) gives

$$u_y \gamma'(u_y) = \frac{c\mu^2 - \mu'}{\alpha''}.$$

Since u, u_y are independent variables then $u_y \gamma'(u_y) = \kappa$. Here κ is a constant. This contradicts to the condition of Lemma.

In the converse case $\alpha_1 \beta_1 \neq 0$, the equation (129) implies $\beta'(u_x) = -\beta_1/(\alpha_1 u_x)$. This is also a contradiction.

(i.2) If (121) holds, then the equation (115) takes the form

$$\mu_i\left(\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)}\right) = 1, \quad \text{for all} \quad i.$$

It means that $\mu_i = 1$, $\alpha_i = \alpha_1$, $\beta_i = \beta_1$ for all i. Thus, we have

$$\beta'(u_x) = \frac{\beta_1}{1 - \alpha_1 u_x}. \tag{131}$$

After substituting (131) in (116), we get that this case is not realized.

(i.3) By rewriting (127), (128) under the case (122), we obtain

$$\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)} = \beta(u_x) + \gamma_1, \quad \mu_i\left(\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)}\right) = \beta(u_x) + \gamma_i.$$

Multiplying the first equation by μ_i and subtracting the second one, we obtain

$$\beta(u_x)(\mu_i - 1) + \gamma_1\mu_i - \gamma_i = 0.$$

Therefore $\mu_i = 1$, $\gamma_i = \gamma_1$ and $\alpha_i = \alpha_1$, $\beta_i = \beta_1$ for all i. Thus, we have

$$\beta' = \frac{\beta_1}{\beta(u_x) - \alpha_1 u_x + \gamma_1}.$$

We substitute this equation in (117) and obtain a contradiction to the condition of Lemma.

(i.4) If (123) holds, then the expressions (127), (128) take the forms

$$\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)} = (\beta + \gamma_1)\exp(\beta + \gamma_1), \quad \mu_i\left(\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)}\right) = (\beta + \gamma_i)\exp(\beta + \gamma_i).$$

Therefore

$$(\beta(\exp\gamma_i - \mu_i\exp\gamma_1) + \gamma_i\exp\gamma_i - \mu_i\gamma_1\exp\gamma_1)\exp\beta = 0.$$

It follows that

$$\exp\gamma_i - \mu_i\exp\gamma_1 = 0, \quad \gamma_i\exp\gamma_i - \mu_i\gamma_1\exp\gamma_1 = 0,$$

and $\gamma_i\exp\gamma_i - \gamma_1\exp\gamma_i = 0$ or $\gamma_i = \gamma_1$ for all i. It means that $\alpha(u) + \gamma(u_y) = \gamma_1$, where γ_1 is a constant. This contradicts to $\gamma_{u_y} \neq 0$.

(i.5) If (124) satisfied, then

$$\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)} = \exp(\beta + \gamma_1). \tag{132}$$

Note that $\beta_1 \neq 0$, otherwise $(\beta' u_x)' = 0$. After replacement $\beta + \gamma_1 \to \beta$, the equation (132) takes the form

$$\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)} = \exp\beta. \tag{133}$$

Substituting (133) in (116), we obtain that $c = -\alpha_1/\beta_1$, $c_1 = -1 - c$. Using (133) in (117) yields to

$$\frac{1}{\beta'}\exp\beta\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(c_1 + cu_y\gamma' + c_2) + \mu' u_y\right) + u_x\left(-\frac{c\mu^2}{\gamma'} + \alpha'' u_y + \frac{\mu'}{\gamma'}\right) -$$
$$-\frac{\alpha_1}{\beta_1}u_x\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(c_1 + cu_y\gamma' + c_2) + \mu' u_y\right) = \exp(\alpha+\gamma)\exp\beta.$$

Since $(\beta' u_x)' \neq 0$, we have that $\exp\beta$, u_x are linearly independent and

$$\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(c_1 + cu_y\gamma' + c_2) + \mu' u_y = \exp(\alpha+\gamma)\beta_1,$$
$$-\frac{\alpha_1}{\beta_1}\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(c_1 + cu_y\gamma' + c_2) + \mu' u_y\right) + \left(-\frac{c\mu^2}{\gamma'} + \alpha'' u_y + \frac{\mu'}{\gamma'}\right) = 0.$$

Thus, we have

$$\begin{gathered}
u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v = \beta(u_x) + \gamma(u_y) + \alpha(u), \quad v_{xy} = \exp v, \\
\alpha_1 u_x + \frac{\beta_1}{\beta'} = \exp\beta, \quad \frac{\beta''}{\beta'^2} = cu_x\beta' + c_1, \quad c_1 = -1 - c, \quad c\beta_1 = -\alpha_1, \\
\frac{\gamma''}{\gamma'^2} = cu_y\gamma' + c_2, \quad \frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(cu_y\gamma' + c_1 + c_2) + \mu' u_y = \exp(\alpha+\gamma)\beta, \\
-\alpha_1\exp(\alpha+\gamma) + \alpha'' u_y + \frac{\mu' - c\mu^2}{\gamma'} = 0.
\end{gathered} \tag{134}$$

(i.6) If (125) holds, then the expressions (127), (128) give

$$\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)} = \exp(\beta + \gamma_1) + 1, \quad \mu_i\left(\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)}\right) = \exp(\beta + \gamma_i) + 1.$$

We multiply the first equation by μ_i and subtract the second one. We obtain

$$\exp\beta\,(\mu_i\exp\gamma_1 - \exp\gamma_i) + \mu_i - 1 = 0.$$

This implies that $\mu_i = 1$, $\gamma_i = \gamma_1$ for all i. It means that $\alpha(u) + \gamma(u_y) = \gamma_1$, where γ_1 is a constant. So we have a conradiction to the condition $\gamma_{u_y} \neq 0$.

(i.7) If (126) satisfied, then by using (127), (128), we get

$$\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)} = \exp(\beta + \gamma_1) + \exp(\sigma(\beta + \gamma_1)),$$
$$\mu_i\left(\alpha_1 u_x + \frac{\beta_1}{\beta'(u_x)}\right) = \exp(\beta + \gamma_i) + \exp(\sigma(\beta + \gamma_i)).$$

Multiplying the first equation by μ_i and subtracting the second one gives

$$\exp\beta\,(\mu_i\exp\gamma_1-\exp\gamma_i)+\exp(\sigma\beta)\,(\mu_i\exp(\sigma\gamma_1)-\exp(\sigma\gamma_i))=0.$$

Since $\sigma\neq 1$, we have

$$\mu_i\exp\gamma_1=\exp\gamma_i,\quad \mu_i\exp(\sigma\gamma_1)=\exp(\sigma\gamma_i).$$

Therefore $\mu_i\exp(\sigma\gamma_1)(\mu_i^{\sigma-1}-1)=0$. Hence $\mu_i=1$. Moreover $\gamma_i=\gamma_1$ for all i and $\alpha(u)+\gamma(u_y)=\gamma_1$. It contradicts to $\gamma_{u_y}\neq 0$.

(ii) There exist such α_i, β_i, $i=1,2$, that $\begin{vmatrix}\alpha_1 & \beta_1\\ \alpha_2 & \beta_2\end{vmatrix}\neq 0$.

By using (117), fixing different values u, u_y, we get

$$\alpha_1u_x+\frac{\beta_1}{\beta'(u_x)}=F(\beta(u_x)+\gamma_1),\quad \alpha_2u_x+\frac{\beta_2}{\beta'(u_x)}=F(\beta(u_x)+\gamma_2).$$

It is clear that

$$\kappa_1F(\beta+\gamma_1)-\kappa_2F(\beta+\gamma_2)=u_x,\quad \kappa_3F(\beta+\gamma_1)-\kappa_4F(\beta+\gamma_2)=\frac{1}{\beta'}. \tag{135}$$

Here

$$\kappa_1=\frac{\beta_2}{\alpha_1\beta_2-\alpha_2\beta_1},\quad \kappa_2=\frac{\beta_1}{\alpha_1\beta_2-\alpha_2\beta_1},\quad \kappa_3=\frac{\alpha_2}{\beta_1\alpha_2-\beta_2\alpha_1},\quad \kappa_4=\frac{\alpha_1}{\beta_1\alpha_2-\beta_2\alpha_1}.$$

(ii.1) If (120) holds, then (135) gives $u_x=0$. Thus, this case is not realized.

(ii.2) If (121) holds, then (135) implies that u_x is a constant.

(ii.3) Let us consider (122). We rewrite (135) and obtain

$$\kappa_1(\beta+\gamma_1)-\kappa_2(\beta+\gamma_2)=u_x,\quad \kappa_3(\beta+\gamma_1)-\kappa_4(\beta+\gamma_2)=\frac{1}{\beta'}.$$

Differentiating the first equation, we get

$$\beta'(\kappa_1-\kappa_2)=1.$$

We substitute this formula in the second equation with noting that $\kappa_1-\kappa_2\neq 0$ by the condition (**ii**), we obtain

$$\beta(\kappa_3-\kappa_4)+\gamma_1\kappa_3-\gamma_2\kappa_4=\kappa_1-\kappa_2.$$

Since $\kappa_3 - \kappa_4 \neq 0$, we have that β is a constant. This contradicts to $\beta_{u_x} \neq 0$.

(ii.4) Let us consider (123). Rewriting (135), we get

$$\begin{aligned} \kappa_1(\beta+\gamma_1)\exp(\beta+\gamma_1) - \kappa_2(\beta+\gamma_2)\exp(\beta+\gamma_2) &= u_x, \\ \kappa_3(\beta+\gamma_1)\exp(\beta+\gamma_1) - \kappa_4(\beta+\gamma_2)\exp(\beta+\gamma_2) &= \frac{1}{\beta'}. \end{aligned}$$

In other words

$$u_x = A\beta\exp\beta + B\exp\beta. \tag{136}$$

Substituting (136) in (135), we obtain

$$\begin{gathered} (A+B)\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(c_1 + cu_y\gamma' + c_2) + \mu' u_y\right) + \\ +B\left(\alpha'' u_y + \frac{\mu' - c\mu^2}{\gamma'}\right) = (\alpha+\gamma)\exp(\alpha+\gamma), \\ A\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(c_1 + cu_y\gamma' + c_2) + \mu' u_y + \alpha'' u_y + \frac{\mu' - c\mu^2}{\gamma'}\right) = \exp(\alpha+\gamma). \end{gathered}$$

Then we insert (136) in (116) and get $c = 1$, $c_1 = -2$. Thus, we have

$$\begin{gathered} u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v = \alpha(u) + \beta(u_x) + \gamma(u_y), \quad v_{xy} = v\exp v, \\ (A+B)\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(u_y\gamma' - 2 + c_2) + \mu' u_y\right) + \\ +B\left(\alpha'' u_y + \frac{\mu' - \mu^2}{\gamma'}\right) = (\alpha+\gamma)\exp(\alpha+\gamma), \\ A\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(u_y\gamma' - 2 + c_2) + \mu' u_y + \alpha'' u_y + \frac{\mu' - \mu^2}{\gamma'}\right) = \exp(\alpha+\gamma), \\ u_x = A\beta\exp\beta + B\exp\beta, \quad \frac{\beta''}{\beta'^2} = u_x\beta' - 2, \frac{\gamma''}{\gamma'^2} = u_y\gamma' + c_2. \end{gathered} \tag{137}$$

(ii.5) The case (124) yields to equations (134).

(ii.6) Let us rewrite (135) by using (125).

$$\begin{aligned} \kappa_1\left(\exp(\beta+\gamma_1) + 1\right) - \kappa_2\left(\exp(\beta+\gamma_2) + 1\right) &= u_x, \\ \kappa_3\left(\exp(\beta+\gamma_1) + 1\right) - \kappa_4\left(\exp(\beta+\gamma_2) + 1\right) &= \frac{1}{\beta'}. \end{aligned}$$

Differentiating the first equation and substituting in the second one, we get

$$\exp\beta(\kappa_3\exp\gamma_1 - \kappa_4\exp\gamma_2 - \kappa_1\exp\gamma_1 + \kappa_2\exp\gamma_2) + \kappa_3 - \kappa_4 = 0.$$

If the expression in the bracket is equal to zero, then $\kappa_3 - \kappa_4 = 0$. In this case β is a constant. This contradicts to $\beta_{u_x} \neq 0$.

(ii.7) If (126) holds, then (135) takes the form

$$\begin{aligned} &\exp\beta(\kappa_1\exp\gamma_1 - \kappa_2\exp\gamma_2) + \exp(\sigma\beta)(\kappa_1\exp(\sigma\gamma_1) - \kappa_2\exp(\sigma\gamma_2)) = u_x,\\ &\exp\beta(\kappa_3\exp\gamma_1 - \kappa_4\exp\gamma_2) + \exp(\sigma\beta)(\kappa_3\exp(\sigma\gamma_1) - \kappa_4\exp(\sigma\gamma_2)) = \frac{1}{\beta'}. \end{aligned} \tag{138}$$

Substituting (138) in (116), we obtain

$$\begin{aligned} (1 + c + c_1)(\kappa_1\exp\gamma_1 - \kappa_2\exp\gamma_2)\exp\beta + \\ +(\sigma^2 + c + c_1\sigma)(\kappa_1\exp(\sigma\gamma_1) - \kappa_2\exp(\sigma\gamma_2))\exp\sigma\beta = 0. \end{aligned} \tag{139}$$

Substituting (138) in (117), we get

$$\begin{aligned} (\kappa_1\exp\gamma_1 - \kappa_2\exp\gamma_2)\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(cu_y\gamma' + c_1 + c_2) + \right.\\ \left. + \mu'u_y + \frac{\mu' - c\mu^2}{\gamma'} + \alpha''u_y\right) = \exp(\alpha + \gamma),\\ (\kappa_1\exp\sigma\gamma_1 - \kappa_2\exp\sigma\gamma_2)\left(\sigma\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(cu_y\gamma' + c_1 + c_2) + \mu'u_y\right) + \right.\\ \left. \frac{\mu' - c\mu^2}{\gamma'} + \alpha''u_y\right) = \exp\sigma(\alpha + \gamma). \end{aligned} \tag{140}$$

Let us consider (139). If $\kappa_1\exp(\sigma\gamma_1) - \kappa_2\exp(\sigma\gamma_2) = 0$, then (140) implies that $\exp\sigma(\alpha+\gamma) = 0$. Thus (139) gives

$$1 + c + c_1 = 0, \quad \sigma^2 + c_1\sigma + c = 0.$$

It implies that $\sigma = c$, $c_1 = -1 - c$. Thus, denoting $A = \kappa_1\exp\gamma_1 - \kappa_2\exp\gamma_2$, $B = \kappa_1\exp(\sigma\gamma_1) - \kappa_2\exp(\sigma\gamma_2)$, we get

$$\begin{aligned} &u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v = \alpha(u) + \beta(u_x) + \gamma(u_y), \quad v_{xy} = \exp v + \exp(\sigma v),\\ &A\exp\beta + B\exp(\sigma\beta) = u_x, \quad \frac{\beta''}{\beta'^2} = \sigma u_x\beta' - 1 - \sigma, \quad \frac{\gamma''}{\gamma'^2} = \sigma u_y\gamma' + c_2\\ &A\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(\sigma u_y\gamma' + c_2 - 1) + \mu'u_y + \frac{\mu'}{\gamma'} + \alpha''u_y\right) = \exp(\alpha + \gamma),\\ &B\left(\sigma\left(\frac{\alpha'\mu}{\gamma'} - \frac{\mu^2}{\gamma'}(\sigma u_y\gamma' + c_2 - \sigma) + \mu'u_y\right) + \frac{\mu'}{\gamma'} + \alpha''u_y\right) = \exp\sigma(\alpha + \gamma). \end{aligned} \tag{141}$$

Now look at each case individually (120)–(126). We will use a symmetry $\beta(u_x) \leftrightarrow \gamma(u_y)$ in (20), (23).

In the case (124), we obtained (134). Using a symmetry $\beta(u_x) \leftrightarrow \gamma(u_y)$, we get

$$\begin{gathered}
\alpha_2 u_y + \frac{\beta_2}{\gamma'} = \exp\gamma, \quad \frac{\gamma''}{\gamma'^2} = c u_y \gamma' + c_2, \quad c_2 = -1 - c, \quad c\beta_2 = -\alpha_2, \\
\frac{\alpha'\mu}{\beta'} - \frac{\mu^2}{\beta'}(c u_x \beta' + c_1 + c_2) + \mu' u_x = \exp(\alpha+\beta)\beta_2, \\
-\alpha_2 \exp(\alpha+\beta) + \alpha'' u_x + \frac{\mu' - c\mu^2}{\beta'} = 0.
\end{gathered} \tag{142}$$

Substituting γ from (142) in (134) and β from (134) in (142), we obtain

$$\begin{gathered}
\frac{1}{\beta_2}(\exp\gamma - \alpha_2 u_y)(\alpha'\mu + 2\mu^2(1+c)) + \mu' u_y = \exp(\alpha+\gamma)\beta_1, \\
-\alpha_1 \exp(\alpha+\gamma) + \alpha'' u_y + (\mu' - c\mu^2)\frac{1}{\beta_2}(\exp\gamma - \alpha_2 u_y) = 0, \\
\frac{1}{\beta_1}(\exp\beta - \alpha_1 u_x)(\alpha'\mu + 2\mu^2(1+c)) + \mu' u_x = \exp(\alpha+\beta)\beta_2, \\
-\alpha_2 \exp(\alpha+\beta) + \alpha'' u_x + (\mu' - c\mu^2)\frac{1}{\beta_1}(\exp\beta - \alpha_1 u_x) = 0.
\end{gathered}$$

Since $\exp\gamma$ and u_y, $\exp\beta$ and u_x are independent, we get

$$\begin{gathered}
u_{xy} = \frac{\mu(u)}{\beta'(u_x)\gamma'(u_y)}, \quad v = \alpha(u) + \beta(u_x) + \gamma(u_y), \quad v_{xy} = \exp v, \\
\alpha_1 u_x + \frac{\beta_1}{\beta'} = \exp\beta, \quad \alpha_2 u_y + \frac{\beta_2}{\gamma'(u_y)} = \exp\gamma, \quad -\frac{\alpha_1}{\beta_1} = -\frac{\alpha_2}{\beta_2} = c.
\end{gathered}$$

$$\alpha'\mu + 2\mu^2(c+1) = \beta_1\beta_2 \exp\alpha, \quad c\beta_1\beta_2 \exp\alpha + \mu' - c\mu^2 = 0, \quad \alpha'' + c(\mu' - c\mu^2) = 0. \tag{143}$$

Examining the system (143) for compatibility, we obtain equations (112), (113).

In the case (123), we get an inconsistent system of equations for α, μ.

Further, the case (126) gives (141). Using symmetry $\beta(u_x) \leftrightarrow \gamma(u_y)$, we get

$$\begin{gathered}
A_2 \exp\gamma + B_2 \exp\sigma\gamma = u_y, \quad \frac{\gamma''}{\gamma'^2} = \sigma u_y \gamma' + c_2, \quad c_2 = -1 - c, \\
\frac{A_2}{\beta'}\left(\alpha'\mu - \mu^2(c_1 - 1) + \mu'\right) + A_2 u_x\left(-\sigma\mu^2 + \mu' + \alpha''\right) = \exp(\alpha+\beta), \\
\frac{B_2}{\beta'}\left(\sigma(\alpha'\mu - \mu^2(c_2 - \sigma)) + \mu'\right) + u_x B_2\left(\sigma(-\sigma\mu^2 + \mu') + \alpha''\right) = \exp\sigma(\alpha+\beta).
\end{gathered} \tag{144}$$

Substituting β from (141) in (144) gives

$$\begin{gathered}
(A\exp\beta + B\sigma\exp\sigma\beta)A_2(\alpha'\mu + \mu^2(2+\sigma) + \mu') + \\
+(A\exp\beta + B\exp\sigma\beta)A_2(\mu' - \sigma\mu^2 + \alpha'') = \exp(\alpha+\beta),
\end{gathered}$$

$$(A\exp\beta + B\sigma\exp(\sigma\beta))B_2\left(\sigma(\alpha'\mu + \mu^2(1+2\sigma) + \mu')\right) +$$
$$(A\exp\beta + B\exp\sigma\beta)B_2\left(\sigma(\mu' - \sigma\mu^2) + \alpha''\right) = \exp\sigma(\alpha+\beta).$$

Since $\exp\beta$, $\exp\sigma\beta$ are independent, we get

$$\begin{gathered} AA_2\left(\alpha'\mu + \mu^2(2+\sigma) + 2\mu' - \sigma\mu^2 + \alpha''\right) = \exp\alpha \\ \sigma\alpha'\mu + \sigma(\sigma+1)\mu^2 + (\sigma+1)\mu' + \alpha'' = 0, \\ BB_2\left(\sigma^2\alpha'\mu + 2\sigma^3\mu^2 + 2\sigma\mu' + \alpha''\right) = \exp(\sigma\alpha). \end{gathered} \tag{145}$$

We examine (145) for compatibility. We obtain equations (114), (115). Thus, (32) is reduced to (111)–(115). This completes the proof of Lemma.

IV. CASE 2. $\varphi = \mathrm{c}\ln \mathrm{u_x} + \mathrm{q}(\mathrm{u}, \mathrm{u_y})$.

We have the following statement for this case.

Lemma 6 *Under the condition* (21) *equations* (1), (2) *and* (3) *up to point transformations take the form:*

$$\begin{gathered} u_{xy} = \frac{\mu(u) - q_u(u,u_y)}{q_{u_y}(u,u_y)}u_x, \quad v_{xy} = c_2\exp v, \quad v = \ln u_x + q(u,u_y), \\ \frac{\mu - q_u}{q_{u_y}}\left(\mu - \frac{\mu - q_u}{q_{u_y}^2}q_{u_yu_y} - 2\frac{q_{uu_y}}{q_{u_y}}\right) + \frac{\mu'}{q_{u_y}} - \frac{q_{uu}}{q_{u_y}} + \mu' = c_2\exp q, \quad q_{uu_y} \neq 0. \end{gathered} \tag{146}$$

Proof. We substitute (21) in (12) and get

$$A(u,u_y)q_{u_y}(u,u_y) - q_u(u,u_y)q_{u_y}(u,u_y)u_y + cq_u(u,u_y) = B(u,u_x)\frac{c}{u_x}.$$

Since u_x, u_y are independent, we obtain

$$Aq_{u_y} - q_uq_{u_y}u_y + cq_u = \mu(u), \quad \frac{Bc}{u_x} = \mu(u). \tag{147}$$

Using (147), we get

$$B = \frac{\mu u_x}{c}, \quad A = \frac{\mu + q_uq_{u_y}u_y - cq_u}{q_{u_y}}.$$

We replace A, B, (21) in (9), (10). This gives

$$f = \frac{\mu - cq_u}{cq_{u_y}}u_x. \tag{148}$$

Then we substitute (21), (148) in (13) and obtain

$$u_x\left(\frac{\mu - cq_u}{cq_{u_y}}\left(\frac{\mu}{c} - \frac{\mu - cq_u}{q_{u_y}^2}q_{u_yu_y} - 2c\frac{q_{uu_y}}{q_{u_y}}\right) + \frac{\mu'}{q_{u_y}} - \frac{q_{uu}}{q_{u_y}} + \frac{\mu'}{c}\right) = F(c\ln u_x + q).$$

Differentiating this equation with respect u_x, we get $cF' = F$, Consequently

$$F(z) = c_2 \exp(z/c). \tag{149}$$

Here c_2 is an arbitrary constant. Thus, using (149) we have

$$u_{xy} = \frac{\mu(u) - cq_u(u, u_y)}{cq_{u_y}(u, u_y)} u_x, \quad v = c \ln u_x + q(u, u_y), \quad v_{xy} = c_2 \exp(v/c)$$
$$\frac{\mu - cq_u}{cq_{u_y}} \left(\frac{\mu}{c} - \frac{\mu - cq_u}{q_{u_y}^2} q_{u_y u_y} - 2c \frac{q_{uu_y}}{q_{u_y}} \right) + \frac{\mu'}{q_{u_y}} - c \frac{q_{uu}}{q_{u_y}} + \frac{\mu'}{c} = c_2 \exp(q/c).$$

Further, the transformation $v/c \to v$, $q/c \to q$, $\mu \to c^2\mu$, $c_2/c \to c_2$ reduces these equations to (146). Hence under the condition (21) we have equations (146). This completes the proof.

V. CASE 3. $\varphi = \alpha(\mathbf{u}) + \kappa(\mathbf{u}) \ln \mathbf{u_x} + \mu(\mathbf{u}) \ln \mathbf{u_y}$.

By substituting (22) in (12), we obtain

$$\left(A(u, u_y) - (\kappa'(u) \ln u_x + \mu'(u) \ln u_y + \alpha'(u))u_y\right) \frac{\mu(u)}{u_y} =$$
$$= \left(B(u, u_y) - (\kappa'(u) \ln u_x + \mu'(u) \ln u_y + \alpha'(u))u_x\right) \frac{\kappa(u)}{u_x}.$$

Therefore

$$\frac{B(u, u_x)\kappa(u)}{u_x} + (\kappa'(u) \ln u_x + \alpha'(u)) (\mu(u) - \kappa(u)) =$$
$$\frac{A(u, u_y)\mu(u)}{u_y} - \mu'(u) \ln u_y (\mu(u) - \kappa(u)).$$

Since u_x, u_y are independent, we have

$$\frac{B(u, u_x)\kappa(u)}{u_x} + (\kappa'(u) \ln u_x + \alpha'(u)) (\mu(u) - \kappa(u)) = \lambda(u),$$
$$\frac{A(u, u_y)\mu(u)}{u_y} - \mu'(u) \ln u_y (\mu(u) - \kappa(u)) = \lambda(u).$$

Moreover, this implies that

$$B = \frac{(\lambda - (\mu - \kappa)(\kappa' \ln u_x + \alpha'))\, u_x}{\kappa}, \quad A = \frac{(\lambda + \mu'(\mu - \kappa) \ln u_y)\, u_y}{\mu}.$$

By replacing A, B in (9), (10) we find f as

$$f = \frac{\lambda - \kappa\mu' \ln u_y - \mu\kappa' \ln u_x - \mu\alpha'}{\kappa\mu} u_x u_y. \tag{150}$$

Substituting (22), (142) in (11) and differentiating the relation obtained with respect to u_x, we get $F'\kappa = F$. On the other hand, by differentiating with respect to u_y, we get $F'\mu = F$. Hence

$$\mu(u) = \kappa(u) = c. \tag{151}$$

Thus φ takes the form $\varphi = \alpha(u) + c\ln(u_x u_y)$ and the case (22) is reduced to (20). This completes the proof.

ACKNOWLEDGMENTS

This work is partially supported by the Scientific and Technological Research Council of Turkey (TÜBİTAK) grant 209T062, Russian Foundation for Basic Research (RFBR) (grants 10-01-91222-CT-a, 11-01-97005-Povoljie-a).

REFERENCES

[1] Liouville, J., "Sur l'equation aux différencespartielles $\partial^2 \log \lambda/\partial u \partial v \pm \lambda/(aa^2) = 0$," J. Math. Pure Appl. **18**, 71 (1853).

[2] Bäcklund, A. V., "Einiges über Curven und Flächentransformationen," Lund Universitëts Arsskrift **10**, (1875).

[3] Bäcklund, A. V., "Om Ytor med konstant negativ kröknig," Lund Universitëts Arsskrift **19**, (1883).

[4] Darboux, G., "Leçon sur la théorie Générale des Surfaces," Paris: Gauthier-Villars **2**, (1915).

[5] Bianchi, L., "Ricerche sulle superficie elicoidali e sulle superficie a curvatura costante" Ann. Sc. Norm. Super. Pisa. **2:1**, 285–341 (1879).

[6] Tzitzeica, G., "Sur une nouvelle classe de surface. I, II," C. R. Academy of Sciences, Paris **150**, 955 –1227 (1910).

[7] Ibragimov, N. Kh., Shabat, A.B., "Evolutionary equations with nontrivial Lie–Backlund group," Funkts. Anal. Prilozh. **14:1**, 25–36 (1980).

[8] Svinolupov, S.I., Sokolov, V.V., and Yamilov, R.I., "On Backlünd transformations for integrable evolution equations," Soviet Math. Dokl. **28:1**, 165-168 (1983).

[9] Khabirov, S. V., "Solutions of evolution equations of second order, obtained by differential substitutions," Math. Problem of Hydrodynamics **85**, 146 – 161 (1988).

[10] Khabirov, S. V., "Infinite-parameter families of solutions of nonlinear differential equations," Mat. Sb. **183:11**, 45–54 (1992).

[11] Sokolov, V. V., "On the symmetries of evolution equations," Uspekhi Mat. Nauk **43:5**, 263 (1988).

[12] Goursat, E., "Leçon sur l'integration des equations aux derivees partielles du second ordre a deux variables independantes," Paris: Hermann **2**, (1896).

[13] Zhiber, A.V., Sokolov, V.V., Startsev, S. Ya., "On nonlinear hyperbolic equations, integrable by Darboux," Reports of the Russian Academy of Sciences **343:6**, 746–748 (1995).

[14] Zhiber, A. V., Sokolov, V. V., "On the Darboux integrable hyperbolic equations," Phys. Lett. A **208**, 303–308 (1995).

[15] Anderson, I.M., Kamran, N., "The variational bicomplex for second order scalar partial differential equations in the plane," Duke Math. J. **87:2**, 265–319 (1997).

[16] Anderson, I.M., Juras, M., "Generalized Laplace invariant and the method of Darboux," Duke Math. J. **89:2**, 351–375 (1997).

[17] Zhiber, A.V., Sokolov, V.V., "Exactly integrable hyperbolic equations of Liouville type," Russ. Math. Surv. **56:1**, 63-106 (2001).

[18] Startsev, S.Ya., "Laplace invariants of hyperbolic equations linearizable by a differential substitution," Teoret. Mat. Fiz. **120:2**, 237-247 (1999).

[19] Startsev, S.Ya.,"Hyperbolic equations admitting differential substitutions," Teoret. Mat. Fiz. **127:1**, 63-74 (2001).

[20] Kuznetsova, M.N., "Laplace transformation and nonlinear hyperbolic equations," Ufimsk. Mat. Zh. **1:3**, 87—96 (2009).

[21] Zhiber, A.V., Murtazina, R.D., "Laplace invariants and Characteristic Lie algebras," Problems of Th. and App. Math. Proceedings of the 30-th regional youth conference: Institute of Math. and Mech., UB of the RAS, Ekaterinburg, 118–122 (2008).

[22] Meshkov, A.G., Sokolov, V.V., "Integrable hyperbolic equations of sin-Gordon type," arXiv:0912.5092 (2009).

[23] Borisov, A.B., Zykov, S.A., "The dressing chains of discrete symmetries and proliferation of nonlinear equations," Theor. Math. Phys. **115:2**, 530–541 (1998).

[24] Zhiber, A.V., Shabat A.B., "Klein-Gordon equations with a nontrivial group," Reports of the Academy of Sciences of the USSR. **247:5**, 1102-1107 (1979).

[25] Kuznetsova, M.N., "The Klein-Gordon equation and differential substitutions of the form $v = \varphi(u, u_x)$," The fundemental mathematics and its applications in the natural science. Proceedings of the conference, Bash. State University, Ufa, 76–85 (2010).